

Class X Mathematics


CBSE Board, Set – 3

General Instructions:

- (i) All questions re compulsory.
- (ii) The question paper consists of 31 questions divided into four sections – A, B, C and D.
- (iii) Section A contains 4 questions of 1 mark each, Section B contains 6 questions of 2 marks each, Section C contains 10 questions of 3 marks each and Section D contains 11 questions of 4 marks each.
- (iv) Use of calculators is not permitted.

Section – A

- Q1.** In Fig. 1, PA and PB are tangents to the circle with centre O such that $\angle APB = 50^\circ$. Write the measure of $\angle OAB$.



Sol.1 In ΔAOB , $OA = OB$ (radius)

$$\therefore \angle OAB = \angle OBA$$

$$= x \text{ (let)}$$

In fig. , OAPB is quadrilateral, $\angle OAP$ and $\angle OBP$ is right angle (radius is perpendicular to the tangent at point of contact)

Using angle sum property of quadrilateral,


$$\therefore \angle AOB + \angle OBP + \angle BPA + \angle OAP = 360^\circ$$

$$\angle AOB + 90^\circ + 50^\circ + 90^\circ = 360^\circ$$

$$\Rightarrow \angle AOB + 230^\circ = 360^\circ$$

$$\Rightarrow \angle AOB = 360^\circ - 230^\circ$$

$$= 130^\circ$$



In ΔOAB ,

$$\angle OAB + \angle AOB + \angle OBA = 180^\circ$$

$$\Rightarrow x + 130^\circ + x = 180^\circ$$

$$\Rightarrow 2x + 130^\circ = 180^\circ$$

$$\Rightarrow 2x = 50^\circ$$

$$\Rightarrow x = 25^\circ$$

Hence $\angle OAB = 25^\circ$

Q2. If $x = -\frac{1}{2}$, is a solution of the quadratic equation $3x^2 + 2kx - 3 = 0$, find the value of k .

Sol. 2 Since it is given that $x = -\frac{1}{2}$ is a solution of the given quadratic equation so it must satisfy the given equation.

So putting $x = -\frac{1}{2}$ in equation.

$$3\left(-\frac{1}{2}\right)^2 + 2k\left(-\frac{1}{2}\right) - 3 = 0.$$

$$3\left(\frac{1}{4}\right) + 2k\left(-\frac{1}{2}\right) - 3 = 0.$$

$$\frac{3}{4} - k - 3 = 0.$$

$$-k - 3 = -\frac{3}{4}$$

$$-k = -\frac{3}{4} + 3$$


Hence the value of k is $-\frac{9}{4}$.

Q3. The top of the two towers of height x and y , standing on level ground, subtend angles of 30° and 60° respectively at the centre of the line joining their feet, then find $x : y$.

Sol.3 Let AB and CD be the two towers of height y and x respectively subtending angles of 60° & 30° .

We are required to find the ratio $x:y$.

Let O be the mid-point of the line joining their feet.



$$OB = OC = -BC \quad \dots (i)$$

Now, in ΔAOB

$$\tan 60^\circ = \frac{OB}{OA} = \frac{-BC}{\frac{BC}{2}}$$

$$\tan 60^\circ = \frac{-BC}{(\frac{BC}{2})} \quad \{\text{from (i)}\}$$

$$\Rightarrow (\sqrt{3}/2) BC = y \quad \dots (ii)$$

And in ΔDOC

$$\tan 30^\circ = \frac{OC}{OD} = \frac{BC}{x}$$

$$\tan 30^\circ = \frac{BC}{x} \quad \{\text{from (i)}\}$$

$$\frac{BC}{2\sqrt{3}} = x \quad \dots (iii)$$

$$x : y = \frac{BC}{2\sqrt{3}} : (\frac{\sqrt{3}}{2})BC$$

$$x : y = \frac{2BC\sqrt{3}}{2\sqrt{3}BC} = \frac{1}{\sqrt{3}} : \frac{\sqrt{3}}{2}$$

$$x : y = 1 : 3$$

Q4. A letter of English alphabet is chosen at random. Determine the probability that the chosen letter is a consonant.

Sol.4 There are 26 English alphabets in total, and so total out comes = 26, out of these 26, 5 are vowels .

$$\Rightarrow \text{Number of Consonants} = \text{favorable outcomes} = 26 - 5 = 21.$$

Here the required probability i.e. prob (chosen is a consonant) = $\frac{21}{26}$.


Section - B

Q5. From a point T outside a circle of centre O, tangents TP and TQ are drawn to the circle. Prove that OT is the right bisector of line segment PQ.

Sol.5 Suppose OT intersects PQ at C.

Then, in ΔQTC , ΔPTC

$TP = TQ$ (Tangents from an external point are equal)



$\angle PTC = \angle QTC$ (TP & TQ are equally inclined to OT)

& $TC = TC$ (common)

By SAS criterion of similarity, are have

$\Delta PTC \cong \Delta QTC$.

$\Rightarrow PC = CQ$ and $\angle PCT = \angle QCT$

But $\angle PCT + \angle QCT = 180^\circ$

$\Rightarrow \angle PCT = \angle QCT = 90^\circ$

$\Rightarrow OT \perp PQ$.

Hence OT is the right bisector of line segment PQ.

Q6. Find the middle term of the A.P. 6, 13, 20, ---, 216.

Sol. 6 The given A.P is,

6, 13, 20, 216.

Here $a = 6$, $d = 13 - 6 = 7$.

$a_n = 216$. (nth term)

$\Rightarrow a + (n - 1) d = 216$.

$6 + (n - 1) 7 = 216$

$\Rightarrow 6 + 7n - 7 = 216$

$\Rightarrow 7n - 1 = 216$

$\Rightarrow 7n = 217$

$\Rightarrow n = 31$.

which is odd.

Hence the middle term is $\left(\frac{31}{2}\right)$ term

$= \left(\frac{31}{2}\right)$ term

$= 16^{\text{th}}$ term

$a_{16} = a + 15 d$


$$= 6 + 15 \cdot 7$$

$$= 6 + 105$$

$$= 111$$

Hence the middle term is 111.

- Q7.** In Fig.2, AB is the diameter of a circle with centre O and AT is a tangent. If $\angle AOQ = 58^\circ$, find $\angle ATQ$.



Sol.7 Since AB is the diameter of the circle

So, $\angle AOQ + \angle QOB = 180^\circ$ (linear pair)

$$58^\circ + \angle QOB = 180^\circ$$

$$\angle QOB = 180^\circ - 58^\circ$$

$$= 122^\circ$$

In $\triangle OBQ$,

$OB = OQ$ (radius)

$$\angle OBQ = \angle OQB = x$$

$$\angle BOQ + \angle OBQ + \angle OQB = 180^\circ$$


$$122^\circ + x + x = 180^\circ$$

$$2x = 58^\circ$$

$$\Rightarrow x = 29^\circ$$

$$\angle OQT = 180^\circ - \angle OQB$$

$$= 180^\circ - 29^\circ$$



$$\angle OQT = 151^\circ$$

AOQT is quadrilateral

$$\angle D = 58^\circ, \quad \angle BAT = 90^\circ \quad \angle OQT = 151^\circ$$

$$\begin{aligned} \text{So, } \angle ATQ &= 260^\circ - (58^\circ + 90^\circ + 151^\circ) \\ &= 360^\circ - 299^\circ \end{aligned}$$

$$\angle ATQ = 61^\circ$$

$$\Rightarrow \angle ATQ = 61^\circ$$

Q8. If A (5, 2), B (2,-2) and C (-2, t) are the vertices of a right angled triangle with $\angle B = 90^\circ$. Then find the value of t.

Sol.8 A (5,2), B (2,-2), C(-2, t) are vertices of right angles triangle

$$\angle B = 90^\circ$$

Using distance formula

$$AB = \sqrt{(5-2)^2 + (2+2)^2}$$

$$BC = \sqrt{(2+2)^2 + (-2-t)^2}$$

$$AC = \sqrt{(5+2)^2 + (2+t)^2}$$

Now, using Pythagoras theorem in ΔABC , we have

$$AB^2 + BC^2 = AC^2.$$

$$(3^2 + 4^2) + (4^2 + (2+t)^2) = T^2 + (2-t)^2$$

$$\Rightarrow 9 + 16 + 16 + 4 + t^2 + 4t = 49 + 4 + t^2 - 4t$$


$$\Rightarrow 41 + 4 + 4t = 53 - 4t$$

$$\Rightarrow 8t = 53 - 4t$$

$$\Rightarrow 8t = 8$$


$$\Rightarrow t = 1$$

Hence the value & t is 1.



Q9. Find the ratio in which the point P(-, -) and B (2,-5).

Sol.9



Let us assume that the point P divides AB in the ratio k : 1.

Then using section formula, we have

$$P = \left(\frac{-}{-}, \frac{-}{-} \right)$$

$$\left(-, - \right) = \left(\frac{-}{-}, \frac{-}{-} \right)$$

$$- = \frac{-}{-}$$

and $-\frac{-}{-} = \frac{-}{-}$

$$- = \frac{-}{-}$$

$$3(k + 1) = (2k + -) 4.$$

$$3k + 3 = 8k + 2.$$

$$5k = 1. \quad k = -$$

Hence the required ratio is k = 1 i.e. - = 1 or 1: 5.

Q10. Solve the following quadratic equation for x:

$$9x^2 - 6b^2x - (a^4 - b^4) = 0$$

Sol.10 $9x^2 - 6b^2x + b^4 - a^4$

$$(3x)^2 - 2(3x)(b^2) + (b^2)^2 - a^4$$

$$(3x - b^2)^2 - (a^2)^2 = 0$$

$$(3x - b^2 + a^2)(3x - b^2 - a^2) = 0$$

$$x = \text{---} \quad x = \text{---}$$

Section -C

Q11. In Fig. 3, APB and AQO are semicircles, and $AO = OB$. If the perimeter of the figure is 40 cm, find the area of the shaded region. [User $\pi = \text{---}$].




Figure 3

Sol.11 $AO = OB$

Perimeter of figure

$$= \pi r + \pi R + R$$

$$r = \text{---}$$

$$\text{Perimeter} = \text{---} + \pi R + R = R(\text{---} + 1)$$

$$40 = R(\text{---} + 1)$$

$$R = \frac{\text{---}}{\text{---}} = \text{---} = \frac{\text{---}}{\text{---}} = 7$$

$$R = 7 \text{ cm. } r = \text{---} = 3.5 \text{ cm}$$


$$\text{Area of shaded region} = \text{---} + \text{---}$$

$$= \text{---} + \text{---}$$

$$= \text{---} + \text{---}$$

$$= \text{---} + \text{---}$$

$$= \text{---} \cdot 77 = 1.25 \times 77 = 96.25 \text{ cm}^2$$



Q12. A solid wooden toy is in the form of a hemisphere surmounted by a cone of same radius. The radius of hemisphere is 3.5 cm and the total wood used in the making of toy is $166\frac{2}{3}$ cm³. Find the height of the toy. Also, find the cost of painting the hemispherical part of the toy at the rate of Rs. 10 per cm². [Use $\pi = \frac{22}{7}$]

Sol. 12 $R = 3.5$ cm

$$V = \frac{2}{3}\pi R^2 H$$

$$\text{Height of cone} = H$$

$$\text{Volume} = \frac{2}{3}\pi R^2 H + \frac{4}{3}\pi R^3$$

$$166\frac{2}{3} = \frac{2}{3}\pi R^2 H + \frac{4}{3}\pi R^3$$


$$166\frac{2}{3} = \frac{2}{3} \times \frac{22}{7} \times 3.5^2 H + \frac{4}{3} \times \frac{22}{7} \times 3.5^3$$

$$6 = H$$

$$H = 6$$
 cm

$$\begin{aligned} \text{Surface area of hemispherical part to painted} &= 2\pi R^2 = 2 \times \frac{22}{7} \times 3.5 \times 3.5 \\ &= 22 \times 3.5 = 77 \text{ cm}^2 \end{aligned}$$

$$\text{Total Cost} = 10 \times 77 = \text{Rs. } 770$$



Q13. In Fig. 4, from the top of a solid cone of height 12 cm and base radius 6 cm, a cone of height 4 cm is removed by a plane parallel to the base. Find the total surface area of the remaining solid. (User $\pi = \frac{22}{7}$ and $\sqrt{5} = 2.236$)

Sol. 13 Using similarity

$$\frac{r}{R} = \frac{h}{H}$$

$$\frac{r}{6} = \frac{h}{12}$$

$$r = 2 \text{ cm}$$

$$r = 2 \text{ cm} \quad (H - h) = 8$$

$$R = 6 \text{ cm}$$

$$l = \sqrt{r^2 + h^2} = \sqrt{4 + 16} = 2\sqrt{5}$$





Figure 4

-- = --

$$L = \frac{2\sqrt{5}}{1} = 6\sqrt{5}$$

$$\begin{aligned} \text{TSA} &= \pi r^2 + \pi R^2 + \pi(L - l)(r + R) \\ &= \pi [4 + 36 + 4\sqrt{5} \times 8] \\ &= 8\pi [5 + 4\sqrt{5}] = 350.592 \text{ cm}^2 \end{aligned}$$

Q14. In Fig. 5, from a cuboidal solid metallic block, of dimensions 15 cm × 10 cm × 5 cm, a cylindrical hole of diameter 7 cm is drilled out. Find the surface area of the remaining block. [Use $\pi = \frac{22}{7}$]




Figure 5

Sol. 14 Surface area of remaining block = Surface area of block - 2 area of circle + C.S.A. of cylinder formed

$$\begin{aligned} &= 2(15 \times 10 + 10 \times 5 + 5 \times 15) - 2 \times \frac{\pi \times 7^2}{4} + 2 \times \pi \times 7 \times 5 \\ &= 2[275] - 77 + 110 \\ &= 583 \text{ cm}^2 \end{aligned}$$

Q15. The angle of elevation of the top of a building from the foot of the tower is 30° and the angle of elevation of the top of the tower from the foot of the building is 45° . If the tower is 30 m high, find the height of the building.

Sol.15 AB → Tower

CD → Building

In ΔABC

$$\tan 45 = \frac{AB}{BC}$$


$$BC = AB$$

$$BC = 30$$

In ΔBDC

$$\tan 30 = \frac{CD}{BC}$$

$$CD = \frac{30}{\sqrt{3}} = 10\sqrt{3} \text{ m.}$$



Q16. In Fig. 6, find the area of the shaded region [Use $\pi = 3.14$]




Figure 6

Sol.16 ABCD will be square with side = diameter of semi-circle

$$\text{Radius} + \text{side} + \text{radius} = 14 - 3 - 3$$

$$r + 2r + r = 8$$

$$r = 2 \text{ cm}$$


$$\text{Area of unshaded figure} = 4^2 + 4 \text{ —}$$

$$= 16 + 25 \cdot 12$$

$$= 41 \cdot 12 \text{ cm}^2$$

$$\text{Area of shaded figure} = 14^2 - 41 \cdot 12$$

$$= 154 \cdot 88 \text{ cm}^2$$



Q17. Find that non-zero value of k , for which the quadratic equation $kx^2 + 1 - 2(k-1)x + x^2 = 0$ has equal roots. Hence find the roots of the equation.

Sol.17 $kx^2 + 1 - 2(k-1)x + x^2 = 0$

$$x^2(k+1) - 2(k-1)x + 1 = 0$$

For equal roots

$$D = 0$$

$$b^2 - 4ac = 0$$

$$4(k - 1)^2 - 4(1)(k + 1) = 0$$

$$4(k^2 + 1 - 2k) - 4k - 4 = 0$$

$$4k^2 + 4 - 8k - 4k - 4 = 0$$

$$4k^2 - 8k - 4k = 0$$

$$k[4k - 12] = 0$$

$$k = 0, k = 3$$

Since non-zero value is required

$$k = 3(\text{Ans.})$$

Q18. All red face cards are removed from a pack of playing cards. The remaining cards were well shuffled and then a card is drawn at random from them. Find the probability that the drawn card is.

Sol.18 6 Red face cards are there remaining cards = 46

(i) Red cards remaining = $26 - 6$

$$= 20$$

Probability = $\frac{6}{20} = \frac{3}{10}$

(ii) Face cards remaining = 6

Probability = $\frac{6}{46} = \frac{3}{23}$

(iii) Card of clubs = 13, Probability = $\frac{13}{46}$

Q19. Find the area of the triangle PQR with Q (3,2) and the mid-points of the side through Q being (2,-1) and (1,2).


Sol.19 Using mid-point formula $\frac{x_1 + x_2}{2} = 2$ $\frac{y_1 + y_2}{2} = -1$

$$x = 1 \quad y = -4$$

$$x = 1 \quad y = 1$$

$$a = -1 \quad b = 0$$

$$P(1, -4)$$



$$R (-1, 0)$$

$$Q (3, 2)$$

$$\text{Area of } \Delta = | - [1 (0 - 2) + (-1) (2 + 4) + 3 (-4 - 0)] |$$

$$- | -(-2 - 6 - 12) |$$

$$= 10 \text{ sq units}$$

Q20. If S_n denotes the sum of first n terms of an A.P., prove that $S_{10} = 3|S_{20} - S_{10}|$

Sol.20 $S_{30} = 3 (S_{20} - S_{10})$

$$S_n = - (2a + (n - 1)d)$$

$$S_{30} = - (2a + 29d) = 15 (2a + 29d) \dots\dots (1)$$

$$3 (S_{20} - S_{10}) = 3 [- (2a + 19d) - - (2a + 9d)]$$

$$= 3 [10a + 145 d]$$

$$= 15 [(2a + 29d)] \dots\dots (2)$$

From (1) and (2)

$$S_{30} = 3 (S_{20} - S_{10})$$


Section - D

Q21. Prove that the tangent at any point of a circle is perpendicular to the radius through the point of contact.

Sol. 21 Given : A circle $C (O, r)$ and a tangent

AB at a point P

To prove : $OP \perp AB$



Construction: Take any point Q, other than P, on the tangent AB. Join OQ suppose OQ meets the circle at R.

Proof : -

We know that among all line segment joining point O to a point on AB, the shortest one is perpendicular to AB.

Hence to prove $OP \perp AB$, we shall first prove that OP is shorter than any other segment joining O to any point of AB .

Clearly, $OP = OR$ (radius)

Now, $OQ = OR + RQ$

$$OQ > OR$$

$$OQ > OP \quad (\because OP = OR)$$

$$OP < OQ$$

Thus, OP is shorter than any other segment joining O to any point of AB

Hence $OP \perp AB$.

Hence the tangent at any point of a circle is perpendicular to the radius through point of contact.

- Q22.** In Fig.7, tangents PQ and PR are drawn from an external point P to a circle with centre O , such that $\angle RPQ = 30^\circ$. A chord RS is drawn parallel to the tangent PQ . Find $\angle RQS$.




Figure 7


Sol. 22 Since, $PQ = PR$

$$\angle PQR = \angle QPR = \frac{180^\circ - 30^\circ}{2} = 75^\circ$$

Now, $\angle RSQ = \angle RQP = 75^\circ$ (because angle in equal segment)

Also, $\angle SQP = 180 - \angle RSQ$ (internal angles)

$$m \angle SQP = 105 \therefore \angle RQS = 30^\circ$$



- Q23.** From a point P on the ground the angle of elevation of the top of a tower is 30° and that of the top of a flag staff fixed on the top of the tower, is 60° . If the length of the flag staff is 5 m, find the height of the tower.

Sol.23 In Δ CBP BC is tower \rightarrow H

$\tan 30 = \frac{h}{AC}$ AC is flag \rightarrow h

$$\frac{1}{\sqrt{3}} = \frac{h}{BP}$$

$$BP = H\sqrt{3} \quad \dots\dots(1)$$

$\tan 60 = \frac{h}{BP}$

$$BP = \frac{h}{\sqrt{3}} \quad \dots\dots(2)$$

From (1) and (2)

$$\frac{h}{\sqrt{3}} = \sqrt{3}H \quad \Rightarrow H + h = 3H$$

$$h = 2H$$

Length of flag staff $h = 5\text{m} = 2H$

Length of tower = 2.5m

Q24. Ramkali required Rs. 2500 after 12 weeks to send her daughter to school. She saved Rs. 100 in the first week and increased her weekly saving by Rs. 20 every week. Find whether she will be able to send her daughter to school after 12 weeks.

What value is generated in the above situation?

Sol.24 First week = 100

Second week = 120

Third week = 140

So it will form A.P. with 12 terms

$$a = 100, d = 20$$

$$= \frac{12}{2} [2 \times 100 + (12 - 1)20]$$

$$= 6[200 + 220]$$

$$= 6 \times 440$$

$$= \text{Rs. } 2640 > \text{Rs. } 2500$$

Yes she will be able to send money.

Q25. The numerator of a fraction is 3 less than its denominator. If 2 is added to both the numerator and the denominator, then the sum of the new fraction and original fraction is $\frac{1}{2}$. Find the original fraction.

Sol.25 Let Denominator be x

Numerator $x - 3$

Fraction = $\frac{x-3}{x}$

New fraction = $\frac{x-1}{x+2}$

ATQ

$\frac{x-3}{x} + \frac{x-1}{x+2} = \frac{1}{2}$

$$\frac{(x-3)(x+2) + (x-1)x}{x(x+2)} = \frac{1}{2}$$

$$2(x-3)(x+2) + 2(x-1)x = x(x+2)$$

$$40x^2 - 40x - 120 = 29x^2 + 58x$$

$$11x^2 - 98x - 120 = 0$$

$$11x^2 - 110x + 12x - 120 = 0$$

$$11x(x-10) + 12(x-10) = 0$$

$$(11x+12)(x-10) = 0$$

$x = \frac{12}{11}$ Rejected as x is integer/whole

$$x = 10$$

Original Fraction = $\frac{7}{10}$

Q26. Water is flowing at the rate of 2.52 km/h through a cylindrical pipe into a cylindrical tank, the radius of whose base is 40 cm, If the increase in the level of water in the tank, in half an hour is 3.15 m, find the internal diameter of the pipe.

Sol.26 Rate of flowing = 2.52 km/hr

$$= 2.52 \times \frac{1000}{60}$$

$$= 0.7 \text{ m/s}$$

Let r be radius of pipe

$$\text{Volume flowing through pipe} = \pi r^2 \cdot 0.7 \text{ —}$$

$$= 0.7 \times \pi r^2$$

$$= 2.2r^2 \text{ —}$$

$$\text{Time} = 30 \text{ min} = 1800\text{s}$$

$$\text{Volume flown} = 2.2 \times r^2 \times 1800 \text{ m}^3$$

In tank

$$\text{Height} = 3.15 \text{ m}$$

$$\text{Radius} = 0.4 \text{ m}$$

Volume flown from pipe = volume rised in tank

$$2.2 \times r^2 \times 1800 = \pi \cdot 0.4 \times 0.4 \times 3.15$$

$$r^2 = \frac{0.4 \cdot 0.4 \cdot 0.45}{2.2}$$

$$r^2 = \frac{0.4 \cdot 0.4 \cdot 0.45}{2.2}$$

$$r = \frac{0.4}{2} = \frac{0.2}{1} = 2 \text{ cm.}$$

$$\text{Diameter} = 2r = 4 \text{ cm.}$$


Q27. If $A(-4, 8)$, $B(-3, -4)$, $(0, -5)$ and $D(5, 6)$ are the vertices of a quadrilateral ABCD, find its area.

Sol.27 A - $(-4, 8)$

B - $(-3, -4)$

C - $(0, -5)$

D - $(-5, 6)$



$$\text{Area of quadrilateral} = \text{Area ABC} + \text{Area ADC}$$

$$\text{Area ABC} = \frac{1}{2} | -4(-4 + 5) + (-3)(-5 - 8) + 0(8 + 4) |$$

$$= -(-4 + 39)$$

$$= - \text{sq unit}$$

$$\text{Area ADC} = | -(-4 [6 + 5] + 5 (-5 -8) + 0 (8 - 6)) |$$

$$= | -(-44 + 65) |$$

$$= 21 \text{ sq units}$$

$$\text{Area of quad} = 21 + 21 = 42 = 72 \text{ sq units}$$

Q28. A 21 m deep well with diameter 6 m is dug and the earth from digging is evenly spread to form a platform 27 m × 11 m. Find the height of the platform. [Use $\pi = \frac{22}{7}$]

Sol.28 volume of earth dug out = volume of embankment

$$\text{Radius} = 3\text{m} \quad \pi \times 3 \times 3 \times 21 = 27 \times 11 \times h$$

$$h = \frac{21 \times 3 \times \pi}{27 \times 11}$$

$$h = 2\text{m}$$

Q29. A bag contains 25 cards numbered from 1 to 25. A card is drawn at random from the bag. Find the probability that the number on the drawn card is:

(i) Divisible by 3 or 5

(ii) a perfect square number

Sol.29 No. of cards Divisible by 3 or 5 = cards divisible by 5 + cards divisible by 3 - cards divisible by 3 and 5

$$= 5 + 8 - 1$$

$$= 12$$

$$\text{Probability} = \frac{12}{25}$$

$$\text{Perfect square} = 1, 4, 9, 16, 21 \quad \text{Probability} = \frac{5}{25} = \frac{1}{5}$$

Q30. Draw a line segment AB of length 7 cm. Taking A as centre, draw a circle of radius 3cm and talking B as centre, draw another circle of radius 2 cm. Construct tangents to each circle from the centre of the other circle.

Sol.30 Steps:-

(i) Draw the perpendicular bisector of AB mark mid -point of AB = X.


(ii) Now from X draw a circle taking AX = BX as the radius

(iii) mark the points of intersection of S, S₂, S₃, S₄

(iv) Join BS and BS₂

Also AS₃ and AS₄

These are required tangents.



Q31. Solve for x:

$$\frac{1}{x-1} + \frac{1}{x+1} = \frac{2}{x^2-1}; x \neq 1, -1$$

Sol.31 $\frac{1}{x-1} + \frac{1}{x+1} = \frac{2}{x^2-1}$ $\frac{3(x-1)+4(x+1)}{x^2-1} = \frac{2}{x^2-1}$

$$(7x + 1) (4x - 1) = 29 (x^2 - 1)$$

$$28x^2 - 3x - 1 = 29x^2 - 29$$

$$x^2 + 3x - 28 = 0$$

$$x^2 + 7x - 4x - 28 = 0$$

$$(x + 7) (x - 4) = 0$$

$$x = -7$$

$$x = 4$$