

## Model Paper for IIT JEE

### Paper I

#### Objective Questions I [Only one correct option]

##### Q 1.

If  $\sqrt{2} \cos A = \cos B + \cos^3 B$ ,  $\sqrt{2} \sin A = \sin B - \sin^3 B$ . Then,  $|\sin(A - B)|$  is equal to

- (a)  $1/2$
- (b)  $1/3$
- (c)  $2/3$
- (d)  $1/5$

##### Q 2.

If  $\alpha$  is a root  $x^4 = 1$ , with negative principal argument, then

Principal argument of  $\omega$  where  $\omega = \begin{vmatrix} 1 & 1 & 1 \\ \alpha^n & \alpha^{n+1} & \alpha^{n+3} \\ \frac{1}{\alpha^{n+1}} & \frac{1}{\alpha^n} & 0 \end{vmatrix}$  (where  $n \in \mathbb{Z}_+$ ) is

- (a)  $\frac{\pi}{4}$
- (b)  $-\frac{3\pi}{6}$
- (c)  $\frac{\pi}{4}$
- (d)  $-\frac{\pi}{4}$

##### Q 3.

If  $a$  and  $b$  be two perpendicular unit vector such that

$x = b - (a \times x)$ , then  $|x|$  is equal to

- (a) 1
- (b)  $\sqrt{2}$
- (c)  $\frac{1}{\sqrt{2}}$
- (d)  $\sqrt{3}$

**Q 4.**

Number of points on hyperbola  $\frac{x^2}{b^2} - \frac{y^2}{a^2} = 1$  from where mutually perpendicular tangents can be drawn to circle  $x^2 + y^2 = a^2$  is

- (a) 2
- (b) 3
- (c) infinite
- (d) 4

**Q 5.**

If  $x + y + z + w = 5$ , then the least value of  $x^2 \cot 9^\circ + y^2 \cot 27^\circ + z^2 \cot 63^\circ + w^2 \cot 81^\circ$  is

- (a)  $\frac{5}{4}$
- (b)  $\frac{5\sqrt{5}}{4}$
- (c)  $\frac{25}{4}$
- (d)  $\frac{25\sqrt{5}}{4}$

**Q 6.**

If  $\sin^{-1} \sin(\sqrt{1-\alpha}) = \sqrt{1-\alpha}$ , when

- (a)  $-\frac{\pi}{2} \leq \alpha \leq \frac{\pi}{2}$
- (b)  $\alpha > \frac{4-\pi}{4}$
- (c)  $\alpha < 4 - \frac{\pi}{4}$
- (d)  $\frac{4-\pi^2}{4} \leq \alpha \leq 1$

**Q 7.**

From any point A on the circle  $|z - z_1| = R$ , tangents are drawn to the circle  $|z - z_1| = 10$ , which meet the former circle at B, C. If chord BC of the former circle touches the later circle, then R is equal to

- (a) 5
- (b) 20
- (c) 10
- (d) 15

**Objective Questions II [One or more than one correct options]****Q 8.**

If  $f(x) = \int_1^x \frac{1 \ln t}{1+t} dt$ , then

(a)  $f\left(\frac{1}{x}\right) = -\int_1^x \frac{1 \ln t}{t(1+t)} dt$

(b)  $f\left(\frac{1}{x}\right) = \int_1^x \frac{1 \ln t}{t(1+t)} dt$

(c)  $f(x) + f\left(\frac{1}{x}\right) = 0$

(d)  $f(x) + \left(\frac{1}{x}\right) = \frac{1}{2}(1 \ln - x)^2$

**Q 9.**

If both the roots of the equation  $ax^2 + x + - a = 0$  are imaginary and  $c > -1$ , then

- (a)  $3a > 2 + 4c$
- (b)  $3a < 2 + 4c$
- (c)  $c < a$
- (d)  $a > 0$

**Q 10.**

If  $\alpha, \beta_i \in \mathbb{R}$  and

$$\sin^2 \theta_1 =$$

$$\frac{(\cos^2 \alpha_1 + \cos^2 \alpha_2 + \cos^2 \alpha_3)(\sin^2 \beta_1 + \sin^2 \beta_2 + \sin^2 \beta_3)}{(\cos \alpha_1 \sin \beta_1 + \cos \alpha_2 \sin \beta_2 + \cos \alpha_3 \sin \beta_3)^2}$$

$$\cos^2 \theta_2 = \frac{(\sin^2 \alpha_1 + \sin^2 \alpha_2 + \sin^2 \alpha_3)(\cos^2 \beta_1 + \cos^2 \beta_2 + \cos^2 \beta_3)}{[\sin \alpha_1 \cos \beta_1 + \sin \alpha_2 \cos \beta_2 + \sin \alpha_3 \cos \beta_3]^2}$$

Then

(a)  $1 = \sin^4 \theta_1$

(b)  $\cos^4 \theta_2 = 1$

(c)  $\sin^8 \theta_1 + \cos^8 \theta_2 = 2$

(d) Cannot be determined

**Q 11.**

Tangent is drawn at any point  $(x_1, y_1)$  other than vertex on the parabola  $y^2 = 4ax$ . If tangents are drawn from any point on this tangent to the circle  $x^2 + y^2 = a^2$ , such that all the chords of contact pass through a fixed point  $(x^2, y+2)$ , then

(a)  $x_1, a, x^2$  are in GP

(b)  $\frac{y_1}{2}, a, y_2$  are in GP

(c)  $-4, \frac{y_1}{y_2}, \frac{x_1}{x_2}$ , are in GP

(d)  $x_1 x_2 + y_1 y_2 = a^2$

**Passage Based Problem**

**Direction (Q. N. 12 to 14)** If the normal at any point P on the ellipse meets the major axis at G and S, S'

Are the foci of the ellipse. Then,  $\frac{SG}{SP} = \frac{S'G}{S'P} = e$  P is any variable point and A, B are any two fixed points. The

Point F moves such that  $PA + PB = k$  ( $k > AB$ ), then locus of P is the ellipse.

**Q 12.**

If the lines  $x + y = 4$  and  $3x + 4y + 5 = 0$  represents equations of the focal chords intersecting at P on the Ellipse whose centre is origin, then equation of tangent at 'P' of the ellipse is

(a)  $x(5 - 3\sqrt{2}) + y(5 - 4\sqrt{2}) - 20 - 5\sqrt{2} = 0$

(b)  $x(5 - 4\sqrt{2} - y(5 - 3\sqrt{2}) + 4 + 5\sqrt{2} = 0$

(c)  $2x - 3y - 5 = 0$

(d)  $2x + 3y + 5 = 0$

**Q 13.**

S (5, 12), S' (-12, 5) are the foci of an ellipse passing through the origin. Then, the eccentricity of ellipse through the origin. Then, the eccentricity of ellipse

(a)  $\frac{1}{\sqrt{13}}$

(b)  $\frac{1}{\sqrt{5}}$

(c)  $\frac{1}{\sqrt{2}}$

(d)  $\frac{1}{\sqrt{7}}$

**Q 14.**

P is any point on the ellipse whose foci are S, S'. Then, w. r. t.  $\Delta SPS'$

(a) the excentre opposite side SS' lies on tangent at P

(b) the excentre opposite to side S' P lies on tangent at P

(c) the excentre opposite to side P' S lies on tangent at P

(d) None of the above

**Directions (Q. No. 15 to 17)** In the Argand plane  $Z_1, Z_2$  and  $Z_3$  are respectively the vertices of an isosceles triangle ABC with  $AC = BC$  and  $\angle CAB = \theta$ . If  $I (Z_4)$  is the in-centre of triangle, then

**Q 15.**

The value of  $\left(\frac{AB}{IA}\right)^2 \left(\frac{AC}{AB}\right)$  is equal to

(a)  $\frac{(Z_2 - Z_1)(Z_1 - Z_3)}{(Z_4 - Z_1)^2}$

(b)  $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)}$

(c)  $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)^2}$

(d)  $\frac{(Z_2 + Z_1)(Z_3 + Z_1)}{(Z_4 + Z_1)}$

**Q 16.**

The value of  $(Z_4 - Z_1)^2 (1 + \cos \theta) \sec \theta$  is

(a)  $(Z_2 - Z_1) (Z_3 - Z_1)$

(b)  $(Z_2 - Z_1) (Z_3 - Z_1) / Z_4 - Z_1$

(c)  $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)^2}$

(d)  $(Z_2 - Z_1) (Z_3 - Z_1)^2$

**Q 17.**

The value of  $(Z_2 - Z_1)^2 \tan \theta \tan (\theta/2)$  is

(a)  $(Z_1 + Z_2 - 2 Z_3) (Z_1 + Z_2 - 2 Z_4)$

(b)  $(Z_1 + Z_2 - Z_3) (Z_1 + Z_2 - Z_4)$

(c)  $(2 Z_3 - Z_1 - Z_2) (Z_1 + Z_2 - 2 Z_4)$

(d)  $(Z_1 + Z_2 + Z_3) (Z_2 + Z_3 - Z_1)$

**Integer Answer Type Questions****Q 18.**

Let a, b be arbitrary real numbers, Find the smallest natural number b for which the equation

$x^2 + 2(a + b)x + (a - b + 8) = 0$  has unequal real roots for all  $a \in \mathbb{R}$ .

**Q 19.**

If  $x$  and  $y$  are non – zero real numbers satisfying

$$xy(x^2 - y^2) = x^2 + y^2, \text{ find the minimum value of } x^2 + y^2.$$

**Q 20.**

From a point  $P(\alpha, \beta)$ , three real and distinct normals are drawn to the parabola  $y^2 = 8x$ . Let  $A, B$  and  $C$  be a feet of normals. The normals at  $A, B$  and  $C$  cut the  $x$  – axis at  $Q, R$  and  $S$  respectively. The value of  $OQ + OR + OS$  ( $O$  is vertex) must be greater than a positive integer  $k$ , then find the greatest value of  $k$ .

**Q 21.**

If the value of  $\int_0^8 (\sqrt{\cot^{-1}(\cot \pi x)} + \cot^{-1}(\cot \pi \sqrt{x})) dx$

$$= a\sqrt{\pi} + (2a\sqrt{2} - b)\pi, \text{ then find } 3a - b.$$

**Q 22.**

If  $\alpha, \beta$  are two distinct real roots of the equation  $ax^3 + x - 1 - a = 0$ , ( $a \neq 1, 0$ ), none of which is

equal to unity, then the value of  $\lim_{x \rightarrow \frac{1}{\alpha}} \frac{(1+a)x^3 - x^2 - a}{(e^{1-\alpha x} - 1)(x-1)}$  is  $\frac{a^{1-(k\alpha-\beta)}}{\alpha}$ . Find the value of  $kl$ .

**Q 23.**

If  $\alpha, \beta$  be the roots of the equation  $x^2 + ax - \frac{1}{2a^2} = 0$ ,  $a$  being a real parameter, then find the least value of  $[\alpha^4 + \beta^4]$  (where  $[.]$  represents greatest integer function).

**Q 24.**

If the area of the region bounded by the curve  $C: y = \tan x$ , the tangent drawn to  $C$  at  $x = \pi/4$  and

The  $x$  – axis is  $\frac{k}{10} \left( \ln 2 - \frac{1}{2} \right)$ , then find the value of  $k$ .

## Paper II

### Objective Questions I [Only one correct option]

#### Q1.

If the system of equations  $x - ky - z = 0$ ,  $kx - y - z = 0$ ,  $x + y - z = 0$  has a non-zero solution, then the possible values of  $k$  are

- (a) -1, 2
- (b) 0, 1
- (c) 1, 2
- (d) -1, 1

#### Q2.

If  $A$  is a square matrix of order  $n \times n$  such that  $A^2 = A$  and  $I$  is a unit matrix of order  $n \times n$ , then  $(I + A)^n$  is equal to

- (a)  $I + 2^n A$
- (b)  $I + (2^n - 1) A$
- (c)  $I - (2^n - 1) A$
- (d) None of these

#### Q3.

If  $f(x)$  is a non-negative continuous function for all  $x \geq 1$ , such that  $f'(x) \leq pf(x)$  where  $p > 0$  and  $f(1) = 0$ , then  $[f(\sqrt{e}) + f(\sqrt{\pi})]$  is equal to

- (a) 0
- (b) negative
- (c) positive
- (d) None of these



**Q 4.**

In a  $\Delta ABC$  is  $BC = a$ ,  $CA = b$  and  $AB = c$  and  $\Delta$  is the area of triangle. Then, least value of

$$\frac{s^2}{P(A/B)} + \frac{(s-a)^2}{P(\bar{A}/B)} + \frac{(s-b)^2}{P(B/A)} + \frac{(s-c)^2}{P(\bar{B}/A)}$$

is, (where A, B are any two events in a sample space such that

$$P(A), P(B) \neq 0$$

(a)  $4\sqrt{3}\Delta$

(b)  $3\sqrt{3}\Delta$

(c)  $6\sqrt{3}\Delta$

(d)  $12\sqrt{3}\Delta$

**Q 5.**

If  $t_n = \sum_{r=0}^n \frac{1}{\binom{n}{r}^k}$  and  $S_n = \sum_{r=0}^n \frac{r}{\binom{n}{r}^k}$ , where  $k \in \mathbb{Z}_+$ , then  $\cos^{-1}\left(\frac{S_n}{nt_n}\right)$  is

(a)  $\frac{\pi}{6}$

(b)  $\frac{\pi}{3}$

(c)  $\frac{\pi}{4}$

(d)  $\frac{\pi}{2}$

**Q 6.**

Let  $S = \sin\sqrt{2} - \sin\sqrt{3}$  and  $C = \cos\sqrt{2} - \cos\sqrt{3}$  then which of the following is correct?

(a)  $S > 0$  and  $C > 0$

(b)  $S > 0$  and  $C < 0$

(c)  $S < 0$  and  $C > 0$

(d)  $S < 0$  and  $C < 0$

**Q7.**

The sum of all the roots of the equation  $\sin\left(\pi \log_3\left(\frac{1}{x}\right)\right) = 0$  in  $(0, 2\pi)$  is

- (a)  $3/2$
- (b)  $4$
- (c)  $9/2$
- (d)  $13/3$

**Q8.**

If  $e^{(\sin^2 x + \sin^4 x + \sin^6 x + \dots + \infty) \ln 3}$ ,  $x \in \left(0, \frac{\pi}{2}\right)$  satisfies the equation  $t^2 - 28t + 27 = 0$ , then the value of

$(\cos x + \sin x)^{-1}$  equals to

- (a)  $\sqrt{3} - 1$
- (b)  $2(\sqrt{3} - 1)$
- (c)  $\sqrt{3} + 1$
- (d)  $1$

**Objective Question II [One or more than one correct options]**

**Q9.**

Let  $(1 + x^2)^2 (1 + x)^n = \sum_{k=0}^{n+4} a_k x^k$ . If  $a_1, a_2, a_3$  are in AP, then  $n$  is (given that  ${}^n C_r = 0$ , if  $n < r$ )

- (a)  $6$
- (b)  $4$
- (c)  $3$
- (d)  $2$

**Q 10.**

If  $f(x) = \int_0^1 e^{|t-x|} dt$  ( $0 \leq x \leq 1$ ), the maximum value of  $f(x)$  is equal to

- (a)  $\sqrt{e} - 1$
- (b)  $2(e - 1)$
- (c)  $(e - 1)$
- (d)  $2(\sqrt{e} - 1)$

**Q 11.**

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = \frac{\sin(\pi \{x\})}{x^4 + 3x^2 + 7}$ , where  $\{ \}$  is fractional function, then

- (a)  $f$  is injective
- (b)  $f$  is not one - one and non - condition
- (c)  $f$  is a surjective
- (d)  $f$  is a zero function

**Q 12.**

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  be two one - one and onto function, such that they are the mirror images of each other about the line  $y = a$ . If  $h(x) = f(x) + g(x)$ , then  $h(x)$  is

- (a) one - one and onto
- (b) only one - one and not onto
- (c) only onto but not one - one
- (d) None of the above

**Integer Answer Type Questions**

**Q 13.**

Suppose that the side lengths of a triangle are three consecutive integers and one of the Angles is twice another. The number of such triangles is .....

**Q 14.**

Let  $A(2\hat{i} + 3\hat{j} + 5\hat{k})$ ,  $B(-\hat{i}, 3\hat{j} + 2\hat{k})$  and  $C(\lambda\hat{i} + 5\hat{j} + \mu\hat{k})$  are vertices of a triangle and its median through  $A$  is equally inclined to the positive directions of the axes. The value of  $2\lambda - \mu$  is equal to ....

**Q 15.**

The number of integral solutions of  $\sum_{i=1}^3 x_j = 24$ ,  $\sum_{i=1}^3 x_i^2 = 21$  and  $\prod_{i=1}^3 x_i = 440$  is .....

**Q 16.**

If  $f(x + y + z) = f(x) + f(y) + f(z)$  with  $f(1) = 1$  and  $f(2) = 2$  and  $x, y, z \in \mathbb{R}$ , then evaluate

$\lim_{n \rightarrow \infty} \frac{\sum_{r=1}^n (4r)f(3r)}{n^3}$  is equal to .....

**Q 17.**

In  $\Delta ABC$ , if BC is unity,  $\sin \frac{A}{2} = x_1$ ,  $\sin \frac{B}{2} = x_2$ ,  $\cos \frac{A}{2} = x_3$  and  $\cos \frac{B}{2} = x_4$  with  $\left(\frac{x_1}{x_2}\right)^{2007} - \left(\frac{x_3}{x_4}\right)^{2006} = 0$ , then the length of AC is .....

**Q 18.**

$x, y \in \mathbb{R}$ ,  $x^2 + y^2 + xy = 1$ , then the minimum value of  $x^3 y + xy^3 + 4$  is .....

**Match the Columns**

**Q 19.**

Match the statements of Column I with values of Column II.

Column I	Column II
(A) Out of machines, two are faulty, they are tested one by one in a random order till both faulty machines are identified, then the probability that only two test are needed.	(p) 1/2
(B) A dice with 6 faces marked 1, 1, 4, 3, 3, 3, is tossed twice. Find the probability of getting sum 4.	(q) 1/4
(C) $(a_1, b_1, c_1)$ , $(a_2, b_2, c_2)$ and $(a_3, b_3, c_3)$ are direction ratios of three perpendicular lines are direction ratio of line equally inclined to them is given by $k(a_1 + a_2 + a_3)$ , $k(b_1 + b_2 + b_3)$ , $k(c_1 + c_2 + c_3)$ . Then k is given by	(r) 1/3
(d) If three points are lying in a plane what is the probability that a triangle will be formed by joining them.	(s) 1/6

**Q 20.**

**Column I**

**Column II**

(A) Let  $a, b, c$  be three mutually perpendicular vectors with same magnitude. If  $x$  satisfies the relation  $a \times \{(x - b) \times a\} + b \times \{(x - c) \times b\} + c \times \{(x - a) \times c\} = 0$ , then  $x$  is equal to

(p)  $\frac{a+b+c}{3}$

(B) The value of  $\lim_{x \rightarrow \infty} (\sqrt[3]{(x+a)(x+b)(x+c)} - x)$

(q) 1

(C)  $\lim_{x \rightarrow 0} \left( \frac{a^x + b^x + c^x}{3} \right)^{1/x}$  equals to

(r)  $\frac{a+b+c}{2}$

(D) Let  $a, b, c$  are distinct reals satisfying  $a^3 + b^3 + c^3 = 3abc$ . If the quadratic equation  $(a + b + c)x^2 + (a + b + c)x + (c + a - b) = 0$ , then a root of the quadratic equation is

(s)  $\sqrt[3]{abc}$