

## Sequences and Series-Solutions

### SUBJECTIVE PROBLEMS :

#### Sol. 1.

Let the two numbers be a and b, then

$$2ab/a + b = 4 \dots(1); a + b/2 = A; \sqrt{ab} = G$$

$$\text{Also } 2A + G^2 = 27 \Rightarrow a + b + ab = 27 \dots\dots\dots (2)$$

Putting  $ab = 27 - (a + b)$  in eqn. (1). We get

$$54 - 2(a + b)/a + b = 4 \Rightarrow a + b = 9 \text{ then } ab = 27 - 9 = 18$$

Solving the two we get  $a = 6, b = 3$  or  $a = 3, b = 6$ , which are the required numbers.

#### Sol. 2.

Let there be n sides in the polygon.

Then by geometry, sum of all n interior angles of polygon =  $(n - 2) * 180^\circ$

Also the angles are in A. P. with the smallest angle =  $120^\circ$ , common difference =  $5^\circ$

$\therefore$  Sum of all interior angles of polygon

$$= n/2[2 * 120 + (n - 1) * 5]$$

Thus we should have

$$n/2 [2 * 120 + (n - 1) * 5] = (n - 2) * 180$$

$$\Rightarrow n/2 [5n + 235] = (n - 2) * 180$$

$$\Rightarrow 5n^2 + 235n = 360n - 720$$

$$\Rightarrow 5n^2 - 125n + 720 = 0 \Rightarrow n^2 - 25n + 144 = 0$$

$$\Rightarrow (n - 16) (n - 9) = 0 \Rightarrow n = 16, 9$$

Also if  $n = 16$  then  $16^{\text{th}}$  angle =  $120 + 15 * 5 = 195^\circ > 180^\circ$

$\therefore$  not possible. Hence  $n = 9$ .

**Sol. 3.**

$a_1, a_2, \dots, a_n$  are in A. P.  $\forall a_i > 0$

$$\therefore a_1 - a_2 = a_2 - a_3 = \dots = a_{n-1} - a_n = d \text{ (a constant)} \dots \dots \dots (1)$$

Now we have to prove

$$\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} = \frac{n-1}{\sqrt{a_1} + \sqrt{a_n}}$$

$$\text{L. H. S.} = \frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}}$$

$$= \frac{\sqrt{a_1} - \sqrt{a_2}}{a_1 - a_2} + \frac{\sqrt{a_2} - \sqrt{a_3}}{a_2 - a_3} + \dots + \frac{\sqrt{a_{n-1}} - \sqrt{a_n}}{a_{n-1} - a_n}$$

$$= \frac{1}{d} * [\sqrt{a_1} - \sqrt{a_2} + \sqrt{a_2} - \sqrt{a_3} + \dots + \sqrt{a_{n-1}} - \sqrt{a_n}]$$

(Using equation (1))

$$= \frac{1}{d} * [\sqrt{a_1} - \sqrt{a_n}]$$

$$= \frac{a_1 - a_n}{d(\sqrt{a_1} + \sqrt{a_n})} = \frac{(n-1)d}{d(\sqrt{a_1} + \sqrt{a_n})} = \frac{n-1}{\sqrt{a_1} + \sqrt{a_n}}$$

= R H S. Hence Proved.

**Sol. 4.**

If possible let for a G. P.

$$T_p = 27 = AR^{p-1} \dots \dots \dots (1)$$

$$T_q = 8 = AR^{q-1} \dots \dots \dots (2)$$

$$T_r = 12 = AR^{r-1} \dots \dots \dots (3)$$

From (1) and (2)

$$R^{p-q} = 27/8 \Rightarrow R^{p-q} = (3/2)^3 \dots \dots \dots (4)$$

From (2) and (3);

$$R^{q-r} = 8/12 \Rightarrow R^{q-r} = (3/2)^{-1} \dots \dots \dots (5)$$

From (4) and (5)

$$R = 3/2 ; p-q = 3 ; q-r = -1$$

$$p - 2q + r = 4; p, q, r \in \mathbb{N} \quad \dots\dots\dots (6)$$

As there can be infinite natural numbers for p, q, and r to satisfy equation (6)

∴ There can be infinite G. P' s.

**Sol. 5.**

$$2 < a, b, c, < 18 \quad a + b + c = 25 \quad \dots\dots\dots (1)$$

$$2, a, b \text{ are AP} \Rightarrow 2a = b + 2$$

$$\Rightarrow 2a - b = 2 \quad \dots\dots\dots (2)$$

$$b, c, 18 \text{ are in GP} \Rightarrow c^2 = 18b \quad \dots\dots\dots (3)$$

$$\text{From (2)} \Rightarrow a = \frac{b+2}{2}$$

$$(1) \Rightarrow b + \frac{b+2}{2} + c = 25 \Rightarrow 3b = 48 - 2c$$

$$(3) \Rightarrow c^2 = 6(48 - 2c) \Rightarrow c^2 + 12c - 288 = 0$$

$$\Rightarrow c = 12, -24 \text{ (rejected)}$$

$$\Rightarrow a = 5, b = 8, c = 12$$

**Sol. 6.**

Given that a, b, c > 0

We know for +ve numbers A. M. ≥ G. M.

∴ For +ve numbers a, b, c we get

$$a + b + c/3 \geq \sqrt[3]{abc} \quad \dots\dots\dots (1)$$

Also for +ve numbers 1/a, 1/b, 1/c, we get

$$\frac{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}{3} \geq \sqrt[3]{\frac{1}{abc}} \quad \dots\dots\dots (2)$$

Multiplying in eqs (1) and (2) we get

$$\left(\frac{a+b+c}{3}\right) \left(\frac{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}{3}\right) \geq \sqrt[3]{abc} \times \frac{1}{\sqrt[3]{abc}}$$

$$\Rightarrow (a + b + c) \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) \geq 9 \text{ Proved.}$$

**Sol. 7.**

Given that  $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot p_3^{\alpha_3} \dots p_k^{\alpha_k} \dots \dots \dots (1)$

Where  $n \in \mathbb{N}$  and  $p_1, p_2, p_3 \dots p_k$  are distinct

Prime numbers.

Taking log on both sides of eq. (1), we get

$\log n = \alpha_1 \log p_1 + \alpha_2 \log p_2 + \dots + \alpha_k \log p_k \dots \dots \dots (2)$

Since every prime number is such that

$p_i \geq 2$

$\therefore \log_e p_i \geq \log_e 2 \dots \dots \dots (3)$

$\forall i = 1 (1) k$

Using (2) and (3) we get

$\log n \geq \alpha_1 \log 2 + \alpha_2 \log 2 + \alpha_3 \log 2 + \dots + \alpha_k \log 2$

$\Rightarrow \log n \geq (\alpha_1 + \alpha_2 + \dots + \alpha_k) \log 2$

$\Rightarrow \log n \geq k \log 2$  Proved

**Sol. 8.**

The given series is

$\sum_{r=0}^n (-1)^r \cdot {}^n C_r \left[ \frac{1}{2^r} + \frac{3^r}{2^{2r}} + \frac{7^r}{2^{2r}} + \frac{15^r}{2^{4r}} + \dots \text{up to } m \text{ terms} \right]$

$\sum_{r=0}^n (-1)^r \cdot {}^n C_r \left( \frac{1}{2} \right)^r + \left( \frac{3}{4} \right)^r + \left( \frac{7}{8} \right)^r + \left( \frac{15}{16} \right)^r + \dots \text{to } m \text{ terms} ]$

Now,  $\sum_{r=0}^n (-1)^r \cdot {}^n C_r \left( \frac{1}{2} \right)^r = 1 - {}^n C_1 \cdot \frac{1}{2} + {}^n C_2 \cdot \frac{1}{2}^2 - {}^n C_3 \cdot \frac{1}{2}^3 + \dots$

$= \left( 1 - \frac{1}{2} \right)^n = \frac{1}{2^n}$

Similarly,  $\sum_{r=0}^n (-1)^r \cdot {}^n C_r \left( \frac{3}{4} \right)^r = \left( 1 - \frac{3}{4} \right)^n = \frac{1}{4^n}$  etc.

Hence the given series is,  $= \frac{1}{2^n} + \frac{1}{4^n} + \frac{1}{8^n} + \frac{1}{16^n} + \dots \text{to } m \text{ terms}$

$= \frac{\frac{1}{2^n} \left( 1 - \left( \frac{1}{2^n} \right)^m \right)}{1 - \frac{1}{2^n}} [\text{Summing the G. P.}] = \frac{2^{mn} - 1}{2^{mn} (2^n - 1)}$

**Sol. 9.**

Let the three distinct real numbers be  $\alpha/r, \alpha, \alpha r$ .

Since sum of squares of three numbers be  $S^2$

$$\therefore \alpha^2/r^2 + \alpha^2 + \alpha^2 r^2 = S^2$$

$$\text{Or } \alpha^2(1 + r^2 + r^4)/r^2 = S^2 \quad \dots\dots\dots (1)$$

Sum of numbers is  $aS$

$$\therefore \alpha(1 + r + r^2)/r = aS \quad \dots\dots\dots (2)$$

Dividing eq. (1) by the square (2), we get

$$\alpha^2(1 + r^2 + r^4)/r^2 * r^2 / \alpha^2(1 + r + r^2)^2 = S^2/a^2S^2$$

$$(1 + 2r^2 + r^4) - r^2 / (1 + r + r^2)^2 = 1/a^2, (1 + r + r^2)(1 - r + r^2) / (1 + r + r^2)^2 = 1/a^2$$

$$1 - r + r^2 / 1 + r + r^2 = 1/a^2 \Rightarrow a^2r^2 - a^2r + a^2 = 1 + r + r^2$$

$$\Rightarrow (a^2 - 1)r^2 - (a^2 + 1)r + (a^2 - 1) = 0$$

$$\Rightarrow r^2 + (1 + a^2/1 - a^2)r + 1 = 0 \quad \dots\dots\dots (3)$$

For real value of  $r, D \geq 0$

$$\Rightarrow (1 + a^2/1 - a^2)^2 - 4 \geq 0$$

$$\Rightarrow 1 + 2a^2 + a^2 - 4 + 8a^2 - 4^2 \geq 0$$

$$\Rightarrow 3a^4 - 10a^2 + 3 \leq 0 \Rightarrow (3a^2 - 1)(a^2 - 3) \leq 0$$

$$\Rightarrow (a^2 - 1/3)(a^2 - 3) \leq 0$$

Clearly the above inequality holds for

$$1/3 \leq a^2 \leq 3$$

But from eq. (3),  $a \neq 1 \therefore a^2 \in (1/3, 1) \cup (1, 3)$ .

**Sol. 10.**

The given equation is

$$\log_{(2x+3)}(6x^2 + 23x + 21)$$

$$= 4 \log_{(2x+3)} (4x^2 + 12x + 9)$$

$$\Rightarrow \log_{(2x+3)} (6x^2 + 23x + 21)$$

$$+ \log_{(3x+7)} (4x^2 + 12x + 9) = 4$$

$$\Rightarrow \log_{(2x+3)} (2x+3) (3x+7) + \log_{(3x+7)} (2x+3)^2 = 4$$

$$\Rightarrow 1 + \log_{(2x+3)} (3x+7) + 2 \log_{(3x+7)} (2x+3) = 4$$

[ Using  $\log ab = \log a + \log b$  and  $\log a^n = n \log a$  ]

NOTE THIS STEP

$$\Rightarrow \log_{(2x+3)} (3x+7) + 2 / \log_{(2x+3)} (3x+7) = 3 \quad [ \text{Using } \log_{ab} = 1/\log_{ba} ]$$

$$\text{Let } \log_{(2x+3)} (3x+7) = y$$

$$\Rightarrow y + 2/y = 3 \Rightarrow y^2 - 3y + 2 = 0$$

$$\Rightarrow (y-1)(y-2) = 0 \Rightarrow y = 1, 2$$

Substituting the values of  $y$  in (1), we get

$$\Rightarrow \log_{(2x+3)} (3x+7) = 1 \quad \text{and } \log_{(2x+3)} (3x+7) = 2$$

$$\Rightarrow 3x+7 = 2x+3 \quad \text{and } 3x+7 = (2x+3)^2$$

$$\Rightarrow x = -4 \quad \text{and } 4x^2 + 9x + 2 = 0$$

$$\Rightarrow x = -4 \quad \text{and } (x+2)(4x+1) = 0$$

$$\Rightarrow x = -4 \quad \text{and } x = 0, x = -1/4$$

As  $\log_a x$  is defined for  $x > 0$  and  $a > 0$  ( $a \neq 1$ ), the possible value of  $x$  should satisfy all of the following inequalities :

$$\Rightarrow 2x+3 > 0 \quad \text{and } 3x+7 > 0$$

$$\text{Also } 2x+3 \neq 1 \quad \text{and } 3x+7 \neq 1$$

Out of  $x = -4$ ,  $x = -2$  and  $x = -1/4$  only  $x = -1/4$

Satisfies the above inequalities.

So only solutions is  $x = -1/4$ .

**Sol. 11.**

Given that  $\log_3 2, \log_3(2^x - 5), \log_3(2^x - 7/2)$  are in A. P.

$$\Rightarrow 2 \log_3(2^x - 5) = \log_3 2 + \log_3(2^x - 7/2)$$

$$\Rightarrow (2^x - 5)^2 = 2(2^x - 7/2)$$

$$\Rightarrow (2^x)^2 - 10 \cdot 2^x + 25 - 2 \cdot 2^x + 7 = 0$$

$$\Rightarrow (2^x)^2 - 10 \cdot 2^x + 25 - 2 \cdot 2^x + 7 = 0$$

$$\Rightarrow (2^x)^2 - 12 \cdot 2^x + 32 = 0$$

Let  $2^x = y$ , then we get,

$$y^2 - 12y + 32 = 0 \Rightarrow (y - 4)(y - 8) = 0$$

$$\Rightarrow y = 4 \text{ or } 8 \Rightarrow 2^x = 2^2 \text{ or } 2^3 \Rightarrow x = 2 \text{ or } 3$$

But for  $\log_3(2x - 5)$  and  $\log_3(2x - 7/2)$  to be defined

$$2^x - 5 > 0 \text{ and } 2^x - 7/2 > 0$$

$$\Rightarrow 2^x > 5 \text{ and } 2^x > 7/2$$

$$\Rightarrow 2^x > 5$$

$$\Rightarrow x \neq 2 \text{ and therefore } x = 3.$$

**Sol. 12.**

Let a and b be two numbers and  $A_1, A_2, A_3, \dots, A_n$  be n A. M's between a and b

Then a,  $A_1, A_2, \dots, A_n, b$  are in A. P. There are  $(n + 2)$  terms in the series, so that

$$a + (n + 1)d = b \Rightarrow d = \frac{b - a}{n + 1}$$

$$\therefore A_1 = a + d = a + \frac{b - a}{n + 1} = \frac{an + b}{n + 1}$$

$$\therefore \text{ATQ } p = \frac{an + b}{n + 1} \quad \dots (1)$$

The first H. M. between a and b, when nHM's are inserted between a and b can be obtained by replacing a by  $1/a$  and b by  $1/b$  in eq. (1) and then taking its reciprocal.

$$\text{Therefore, } q = \frac{1}{\frac{1}{a} + \frac{1}{b} + \frac{n}{ab}} = \frac{ab}{bn + a}$$

$$\therefore q = \frac{ab}{bn + a} \quad \dots (2)$$

We have to prove that  $q$  cannot lie between  $p$  and  $(n + 1)^2/(n - 1)^2 p$ .

Now,  $n + 1 > n - 1 \Rightarrow n + 1/n - 1 > 1$

$\Rightarrow (n + 1/n - 1)^2 > 1$  or  $p(n + 1/n - 1)^2 > p$

$\Rightarrow p < p(n + 1/n - 1)^2$  ..... (3)

Now to prove the given, we have to show that  $q$  is less than  $p$ .

For this, let,  $p/q = (na + b)(nb + a)/(n + 1)^2 ab$

$\Rightarrow p/q - 1 = n(a^2 + b^2) + ab(n^2 + 1) - (n + 1)^2 ab / (n + 1)^2 ab$

$\Rightarrow p/q - 1 = n(a^2 + b^2 - 2ab) / (n + 1)^2 ab$

$\Rightarrow p/q - 1 = n / (n + 1)^2 (a - b/\sqrt{ab})^2$

$= n / (n + 1)^2 (\sqrt{a/b} - \sqrt{b/a})^2 \Rightarrow p/q - 1 > 0$

$\Rightarrow$  ( provided  $a$  and  $b$  and hence  $p$  and  $q$  are +ve )

$p > q$  ..... (4)

From (3) and (4), we get,

$q < p < (n + 1/n + 1)^2 p$

$\therefore q$  can not lie between  $p$  and  $(n + 1/n + 1)^2 p$ , if  $a$  and  $b$  are +ve numbers.

ALTERNATE SOLUTION:

After getting equations (1) and (2) as in the previous method, substitute  $b = p(n + 1) - an$  [from (1)] in

equation (2) to get  $aq + nq [p(n + 1) - an] = (n + 1)a [p(n + 1) - an]$

$\Rightarrow a^2 n (n + 1) + a [q(1 - n^2) - p(n + 1)^2] + npq (n + 1) = 0$

$\Rightarrow na^2 - [(n + 1)p + (n - 1)q] a + npq = 0$

$\Rightarrow D \geq 0$  (  $\because a$  is real )

$\Rightarrow [n + 1]p + (n - 1)q]^2 - 4n^2 pq \geq 0$

$\Rightarrow (n - 1)^2 q^2 + \{2(n^2 - 1) - 4n^2\} pq + (n + 1)^2 p^2 \geq 0$

$\Rightarrow q^2 - 2n^2 + 1 / (n - 1)^2 pq + (n + 1/n - 1)^2 p^2 \geq 0$

$\Rightarrow [q - p(n + 1/n - 1)^2] [q - p] \geq 0$

[ On factorizing by discriminant method ]  $\Rightarrow q$  can not lie between  $p$  and  $p(n + 1/n - 1)^2$ .

**Sol. 13.**

ATQ we have,

$$S_1 = 1 + \frac{1}{2} + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{3}\right)^3 + \dots \infty$$

$$S_2 = 2 + 2 \cdot \frac{1}{3} + 2 \left(\frac{1}{3}\right)^2 + \dots \infty$$

.....  
.....

$$S_3 = 3 + 3 \cdot \frac{1}{4} + 3 \left(\frac{1}{4}\right)^2 + \dots \infty$$

$$S_n = n + n \frac{1}{n+1} + n \left(\frac{1}{n+1}\right)^2 + \dots \infty$$

$$\Rightarrow S_1 = \frac{1}{1-\frac{1}{2}} = 2 \text{ [using } S_\infty = \frac{a}{1-r}\text{]}$$

$$S_2 = \frac{2}{1-\frac{1}{3}} = 3, \quad S_3 = \frac{3}{1-\frac{1}{4}} = 4,$$

$$S_n = \frac{n}{1-\frac{1}{n+1}} = (n+1)$$

$$\begin{aligned} \therefore S_1^2 + S_2^2 + S_3^2 + \dots + S_{2n-1}^2 \\ = 2^2 + 3^2 + 4^2 + \dots + (n+1)^2 + \dots + (2n)^2 \end{aligned}$$

NOTE THIS STEP:

$$\begin{aligned} \sum_{r=1}^{2n} r^2 - 1 &= \frac{2n(2n+1)(4n+1)}{6} - 1^2 \\ &= \frac{n(2n+1)(4n+1)3}{3} \end{aligned}$$

**Sol. 14.**

Since  $x_1, x_2, x_3$  are in A. P. T. Therefore, let  $x_1 = a - d, x_2 = a$  and  $x_3 = a + d$  and  $x_1, x_2, x_3$  are the roots of

$$x^3 - x^2 + \beta x + \gamma = 0$$

$$\text{We have } \sum \alpha = a - d + a + a + d = 1 \quad \dots (1)$$

$$\sum \alpha \beta = (a - d)a + a(a + d) + (a - d)(a + d) = \beta \quad \dots (2)$$

$$\alpha \beta \gamma = (a - d)a(a + d) = -\gamma \quad \dots (3)$$

From (1), we get,  $3a = 1 \Rightarrow a = 1/3$

From (2), we get  $3a^2 - a^2 = \beta$

$$\Rightarrow 3(1/3)^2 - a^2 = \beta \Rightarrow 1/3 - \beta = d^2$$

(NOTE : In this equation we have two variables  $\beta$  and  $d$  but we have only one equation. So at first sight it looks that this equation cannot be solved but we know that

$d^2 \geq 0 \forall d \in \mathbb{R}$  then ;  $\beta$  can be solved).

$$\Rightarrow 1/3 - \beta \geq 0 \quad \because d^2 \geq 0$$

$$\Rightarrow \beta \leq 1/3 \Rightarrow \beta \in (-\infty, 1/3]$$

From (3),  $a(a^2 - d^2) = -\gamma$

$$\Rightarrow 1/3 (1/9 - d^2) = -\gamma \Rightarrow 1/27 - 1/3d^2 = -\gamma$$

$$\Rightarrow \gamma + 1/27 = 1/3d^2 \Rightarrow \gamma + 1/27 \geq 0$$

$$\Rightarrow \gamma \geq -1/27 \Rightarrow -\gamma \in [-1/27, \infty)$$

Hence  $\beta \in (-\infty, 1/3)$  and  $\gamma \in [-1/27, \infty]$

**Sol. 15.**

Solving the system of equations,  $u + 2v + 3w = 6$ ,

$$4u + 5v + 6w = 12 \text{ and } 6u + 9v = 4$$

We get  $u = -1/3, v = 2/3, w = 5/3$

$$\therefore u + v + w = 2, 1/u + 1/v + 1/w = -9/10$$

Let  $r$  be the common ratio of the G. P.,  $a, b, c, d$ . Then  $b = ar, c = ar^2, d = ar^3$ .

Then the first equation

$$(1/u + 1/v + 1/w) x^2 + [(b - c)^2 + (c - a)^2 + (d - b)^2] x + (u + v + w) = 0$$

Becomes

$$\frac{9}{10}x^2 + [(ar - ar^2)^2 + (ar^2 - a)^2 + (ar^3 - ar)^2] x + 2 = 0$$

$$\text{i.e. } 9x^2 - 10a^2(1 - r)^2 [r^2 + (r + 1)^2 + r^2(r + 1)^2]x - 20 = 0$$

$$\text{i.e. } 9x^2 - 10a^2(1 - r)^2(r^4 + 2r^3 + 3r^2 + 2r + 1)x - 20 = 0$$

$$\text{i.e. } 9x^2 - 10a^2(1 - r)^2(1 + r + r^2)^2 x - 20 = 0,$$

$$\text{i.e. } 9x^2 - 10a^2(1 - r^3)^2 x - 20 = 0 \quad \dots\dots (1)$$

The second equation is

$$20x^2 + 10(a - ar^3)^2 x - 9 = 0$$

i.e.,  $20x^2 + 10a^2 (1 - r^3)^2 x - 9 = 0$  ..... (2)

Since (2) can be obtained by the substitution  $x \rightarrow 1/x$ , equations (1) and (2) have reciprocal roots.

**Sol. 16.**

Let  $a - 3d, a - d, a + d$  and  $a + 3d$  be any four consecutive terms of an A. P. with common difference  $2d$  ∴ Terms of A. P. are integers,  $2d$  is also an integer.

Hence  $p = (2d)^4 + (a - 3d)(a - d)(a + d)(a + 3d)$

$$= 16d^4 + (a^2 - 9d^2)(a^2 - d^2) = a^2 - 5d^2$$

Now,  $a^2 - 5d^2 = a^2 - 9d^2 + 4d^2$

$$= (a - 3d)(a + 3d) + (2d)^2 = \text{some integer}$$

Thus  $p = \text{square of an integer.}$

**Sol 17.**

Given that  $a_1, a_2, \dots, a_n$  are +ve real no's is G. P.

$$\left. \begin{array}{l} a_1 = a \\ a_2 = ar \\ a_3 = ar^2 \\ a_n = ar^{n-1} \end{array} \right\} \text{As } a_1, a_2, \dots, a_n \text{ are +ve } \therefore r > 0$$

$A_n$  is A. M. of  $a_1, a_2, \dots, a_n$

$$\therefore A_n = a_1 + a_2 + \dots + a_n/n = a + ar + \dots + ar^{n-1}/n$$

$$A_n = \frac{a(1-r)^n}{n(1-r)} \dots \dots \dots (1) \text{ (For } r \neq 1)$$

$G_n$  is G. M. of  $a_1, a_2, \dots, a_n$

$$\therefore G_n = \sqrt[n]{a_1 \cdot a_2 \cdot \dots \cdot a_n} = \sqrt[n]{a \cdot ar \cdot ar^2 \cdot \dots \cdot ar^{n-1}}$$

$$= \sqrt[n]{a^n \cdot r^{\frac{n(n-1)}{2}}} = ar^{\frac{(n-1)}{2}}$$

$$G_n = ar^{\frac{(n-1)}{2}} \dots \dots \dots (2) \text{ (} r \neq 1)$$

$H_n$  is H. M. of  $a_1, a_2, \dots, a_n$

$$\begin{aligned} \therefore H_n &= \frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n}} = \frac{n}{\frac{1}{a} + \frac{1}{ar} + \dots + \frac{1}{ar^{n-1}}} \\ &= \frac{n}{\frac{1}{a} \left( \frac{1-r^n}{1-r} \right)} = \frac{n}{\frac{1}{a} \left( \frac{1-r^n}{r^n} \right) \cdot \frac{r}{1-r}} \end{aligned}$$

$$H_n = an r^{n-1} (1-r)/(1-r^n) \quad (r \neq 1) \quad \dots \dots \dots (3)$$

We also observe that

$$A_n H_n = \frac{a(1-r^n)}{n(1-r)} \times \frac{anr^{n-1}(1-r)}{1-r^n} = a^n r^{n-1} = G_n^2$$

$$\therefore A_n H_n = G_n^2 \quad \dots \dots \dots (4)$$

$\therefore$  Now, G. M. of  $G_1, G_2, \dots, G_n$  is

$$G = \sqrt[n]{G_1 G_2 \dots G_n}$$

$$G = \sqrt[n]{\sqrt{A_1 H_1} \sqrt{A_2 H_2} \dots \sqrt{A_n H_n}} \quad [\text{Using (4)}]$$

$$G = (A_1 A_2 \dots A_n H_1 H_2 \dots H_n)^{1/2n} \quad \dots \dots \dots (5)$$

If  $r = 1$  then

$$A_n = G_n = H_n = a$$

$$\text{Also } A_n H_n = G_n^2$$

$\therefore$  For  $r = 1$  also, equation (5) holds.

Hence we get

$$G = (A_1 A_2 \dots A_n H_1 H_2 \dots H_n)^{1/2n}$$

**Sol. 18.**

Clearly  $A_1 + A_2 = a + b$

$$1/H_1 + 1/H_2 = 1/a + 1/b$$

$$\Rightarrow H_1 + H_2/H_1 H_2 = a + b/ab = A_1 + A_2/G_1 G_2$$

$$\Rightarrow G_1 G_2 / H_1 H_2 = A_1 + A_2/H_1 + H_2$$

$$\text{Also } 1/H_1 = 1/a + 1/3 (1/b - 1/a) \Rightarrow H_1 = 3ab/2b + a$$

$$1/H_2 = 1/a + 2/3(1/b - 1/a) \Rightarrow H_2 = 3ab/2a + b$$

$$\Rightarrow A_1 + A_2/H_1 + H_2 = \frac{a+b}{3ab\left(\frac{1}{2b+a} + \frac{1}{2a+b}\right)}$$

$$= \frac{(2b+a)(2a+b)}{9ab}$$

**Sol. 19.**

Given that a, b, c are in A. P.

$$\Rightarrow 2b = a + c \quad \dots\dots (1)$$

And  $a^2, b^2, c^2$  are in H. P.

$$\Rightarrow \frac{1}{b^2} - \frac{1}{a^2} = \frac{1}{c^2} - \frac{1}{b^2}$$

$$\Rightarrow (a-b)(a+b)/b^2a^2 = (b-c)(b+c)/b^2c^2$$

$$\Rightarrow ac^2 + bc^2 = a^2b + a^2c \quad [\because a-b = b-c]$$

$$\Rightarrow ac(c-a) + b(c-a)(c+a) = 0$$

$$\Rightarrow (c-a)(ab+bc+ca) = 0$$

$$\Rightarrow \text{either } c-a = 0 \text{ or } ab+bc+ca = 0$$

$$\Rightarrow \text{either } c = a \text{ or } (a+c)b + ca = 0 \text{ and then from (i) } 2b^2 + ca = 0$$

Either  $a = b = c$  or  $b^2 = a(-c/2)$

i.e. a, b, -c/2 are in G. P. Hence Proved

**Sol. 20.**

$$a_n = 3/4 - (3/4)^2 + (3/4)^3 + \dots\dots + (-1)^{n-1} (3/4)^n$$

$$= \frac{\frac{3}{4}\left(1 - \left(-\frac{3}{4}\right)^n\right)}{1 + \frac{3}{4}} = 3/7 \left(1 - (-3/4)^n\right)$$

$$B_n = 1 - a_n \quad \text{and } b_n > a_n \quad \forall n \geq n_0$$

$$\therefore 1 - a_n > a_n \Rightarrow 2a_n < 1$$

$$\Rightarrow \frac{6}{7} \left[1 - \left(-\frac{3}{4}\right)^n\right] < 1 \Rightarrow -\left(-\frac{3}{4}\right)^n < 1/6$$

$$\Rightarrow (-3)^{n+1} < 2^{2n-1}$$

For n to be even, inequality always holds. For n to be odd, it holds for  $n \geq 7$ .

$\therefore$  The least natural no, for which it holds is 6 (  $\because$  it holds for every even natural no.)