

Sample Question Paper
CLASS: XII
Session: 2021-22
Mathematics (Code-041)
Term - 1

Time Allowed: 90 minutes

Maximum Marks: 40

General Instructions:


1. This question paper contains three sections – A, B and C. Each part is compulsory.
2. Section - A has 20 MCQs, attempt any 16 out of 20.
3. Section - B has 20 MCQs, attempt any 16 out of 20
4. Section - C has 10 MCQs, attempt any 8 out of 10.
5. All questions carry equal marks.
6. There is no negative marking.

SECTION – A

In this section, attempt any 16 questions out of Questions 1 – 20.
 Each Question is of 1 mark weightage.

1.	$\sin \left[\frac{\pi}{3} - \sin^{-1} \left(-\frac{1}{2} \right) \right]$ is equal to:	1				
	<table><tr><td>a) $\frac{1}{2}$</td><td>b) $\frac{1}{3}$</td></tr><tr><td>c) -1</td><td>d) 1</td></tr></table>	a) $\frac{1}{2}$	b) $\frac{1}{3}$	c) -1	d) 1	
a) $\frac{1}{2}$	b) $\frac{1}{3}$					
c) -1	d) 1					
2.	<p>The value of k ($k < 0$) for which the function f defined as</p> $f(x) = \begin{cases} \frac{1 - \cos kx}{x \sin x}, & x \neq 0 \\ \frac{1}{2}, & x = 0 \end{cases}$ <p>is continuous at $x = 0$ is:</p>	1				
	<table><tr><td>a) ± 1</td><td>b) -1</td></tr><tr><td>c) $\pm \frac{1}{2}$</td><td>d) $\frac{1}{2}$</td></tr></table>	a) ± 1	b) -1	c) $\pm \frac{1}{2}$	d) $\frac{1}{2}$	
a) ± 1	b) -1					
c) $\pm \frac{1}{2}$	d) $\frac{1}{2}$					
3.	<p>If $A = [a_{ij}]$ is a square matrix of order 2 such that $a_{ij} = \begin{cases} 1, & \text{when } i \neq j \\ 0, & \text{when } i = j \end{cases}$, then A^2 is:</p>	1				
	<table><tr><td>a) $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$</td><td>b) $\begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix}$</td></tr><tr><td>c) $\begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix}$</td><td>d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$</td></tr></table>	a) $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$	b) $\begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix}$	c) $\begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix}$	d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	
a) $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$	b) $\begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix}$					
c) $\begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix}$	d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$					
4.	<p>Value of k, for which $A = \begin{bmatrix} k & 8 \\ 4 & 2k \end{bmatrix}$ is a singular matrix is:</p>	1				
	<table><tr><td>a) 4</td><td>b) -4</td></tr><tr><td>c) ± 4</td><td>d) 0</td></tr></table>	a) 4	b) -4	c) ± 4	d) 0	
a) 4	b) -4					
c) ± 4	d) 0					

5.	Find the intervals in which the function f given by $f(x) = x^2 - 4x + 6$ is strictly increasing:	1				
<table><tr><td>a) $(-\infty, 2) \cup (2, \infty)$</td><td>b) $(2, \infty)$</td></tr><tr><td>c) $(-\infty, 2)$</td><td>d) $(-\infty, 2] \cup (2, \infty)$</td></tr></table>			a) $(-\infty, 2) \cup (2, \infty)$	b) $(2, \infty)$	c) $(-\infty, 2)$	d) $(-\infty, 2] \cup (2, \infty)$
a) $(-\infty, 2) \cup (2, \infty)$	b) $(2, \infty)$					
c) $(-\infty, 2)$	d) $(-\infty, 2] \cup (2, \infty)$					
6.	Given that A is a square matrix of order 3 and $ A = -4$, then $ \text{adj } A $ is equal to:	1				
<table><tr><td>a) -4</td><td>b) 4</td></tr><tr><td>c) -16</td><td>d) 16</td></tr></table>			a) -4	b) 4	c) -16	d) 16
a) -4	b) 4					
c) -16	d) 16					
7.	A relation R in set $A = \{1, 2, 3\}$ is defined as $R = \{(1, 1), (1, 2), (2, 2), (3, 3)\}$. Which of the following ordered pair in R shall be removed to make it an equivalence relation in A ?	1				
<table><tr><td>a) $(1, 1)$</td><td>b) $(1, 2)$</td></tr><tr><td>c) $(2, 2)$</td><td>d) $(3, 3)$</td></tr></table>			a) $(1, 1)$	b) $(1, 2)$	c) $(2, 2)$	d) $(3, 3)$
a) $(1, 1)$	b) $(1, 2)$					
c) $(2, 2)$	d) $(3, 3)$					
8.	If $\begin{bmatrix} 2a + b & a - 2b \\ 5c - d & 4c + 3d \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 11 & 24 \end{bmatrix}$, then value of $a + b - c + 2d$ is:	1				
<table><tr><td>a) 8</td><td>b) 10</td></tr><tr><td>c) 4</td><td>d) -8</td></tr></table>			a) 8	b) 10	c) 4	d) -8
a) 8	b) 10					
c) 4	d) -8					
9.	The point at which the normal to the curve $y = x + \frac{1}{x}$, $x > 0$ is perpendicular to the line $3x - 4y - 7 = 0$ is:	1				
<table><tr><td>a) $(2, 5/2)$</td><td>b) $(\pm 2, 5/2)$</td></tr><tr><td>c) $(-1/2, 5/2)$</td><td>d) $(1/2, 5/2)$</td></tr></table>			a) $(2, 5/2)$	b) $(\pm 2, 5/2)$	c) $(-1/2, 5/2)$	d) $(1/2, 5/2)$
a) $(2, 5/2)$	b) $(\pm 2, 5/2)$					
c) $(-1/2, 5/2)$	d) $(1/2, 5/2)$					
10.	$\sin(\tan^{-1}x)$, where $ x < 1$, is equal to:	1				
<table><tr><td>a) $\frac{x}{\sqrt{1-x^2}}$</td><td>b) $\frac{1}{\sqrt{1-x^2}}$</td></tr><tr><td>c) $\frac{1}{\sqrt{1+x^2}}$</td><td>d) $\frac{x}{\sqrt{1+x^2}}$</td></tr></table>			a) $\frac{x}{\sqrt{1-x^2}}$	b) $\frac{1}{\sqrt{1-x^2}}$	c) $\frac{1}{\sqrt{1+x^2}}$	d) $\frac{x}{\sqrt{1+x^2}}$
a) $\frac{x}{\sqrt{1-x^2}}$	b) $\frac{1}{\sqrt{1-x^2}}$					
c) $\frac{1}{\sqrt{1+x^2}}$	d) $\frac{x}{\sqrt{1+x^2}}$					
11.	Let the relation R in the set $A = \{x \in \mathbb{Z} : 0 \leq x \leq 12\}$, given by $R = \{(a, b) : a - b \text{ is a multiple of } 4\}$. Then $[1]$, the equivalence class containing 1, is:	1				
<table><tr><td>a) $\{1, 5, 9\}$</td><td>b) $\{0, 1, 2, 5\}$</td></tr><tr><td>c) ϕ</td><td>d) A</td></tr></table>			a) $\{1, 5, 9\}$	b) $\{0, 1, 2, 5\}$	c) ϕ	d) A
a) $\{1, 5, 9\}$	b) $\{0, 1, 2, 5\}$					
c) ϕ	d) A					
12.	If $e^x + e^y = e^{x+y}$, then $\frac{dy}{dx}$ is:	1				
<table><tr><td>a) e^{y-x}</td><td>b) e^{x+y}</td></tr><tr><td>c) $-e^{y-x}$</td><td>d) $2e^{x-y}$</td></tr></table>			a) e^{y-x}	b) e^{x+y}	c) $-e^{y-x}$	d) $2e^{x-y}$
a) e^{y-x}	b) e^{x+y}					
c) $-e^{y-x}$	d) $2e^{x-y}$					

13.	Given that matrices A and B are of order $3 \times n$ and $m \times 5$ respectively, then the order of matrix $C = 5A + 3B$ is:	1				
<table><tr><td>a) 3×5</td><td>b) 5×3</td></tr><tr><td>c) 3×3</td><td>d) 5×5</td></tr></table>			a) 3×5	b) 5×3	c) 3×3	d) 5×5
a) 3×5	b) 5×3					
c) 3×3	d) 5×5					
14.	If $y = 5 \cos x - 3 \sin x$, then $\frac{d^2 y}{dx^2}$ is equal to:	1				
<table><tr><td>a) $-y$</td><td>b) y</td></tr><tr><td>c) $25y$</td><td>d) $9y$</td></tr></table>			a) $-y$	b) y	c) $25y$	d) $9y$
a) $-y$	b) y					
c) $25y$	d) $9y$					
15.	For matrix $A = \begin{bmatrix} 2 & 5 \\ -11 & 7 \end{bmatrix}$, $(adj A)'$ is equal to:	1				
<table><tr><td>a) $\begin{bmatrix} -2 & -5 \\ 11 & -7 \end{bmatrix}$</td><td>b) $\begin{bmatrix} 7 & 5 \\ 11 & 2 \end{bmatrix}$</td></tr><tr><td>c) $\begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix}$</td><td>d) $\begin{bmatrix} 7 & -5 \\ 11 & 2 \end{bmatrix}$</td></tr></table>			a) $\begin{bmatrix} -2 & -5 \\ 11 & -7 \end{bmatrix}$	b) $\begin{bmatrix} 7 & 5 \\ 11 & 2 \end{bmatrix}$	c) $\begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix}$	d) $\begin{bmatrix} 7 & -5 \\ 11 & 2 \end{bmatrix}$
a) $\begin{bmatrix} -2 & -5 \\ 11 & -7 \end{bmatrix}$	b) $\begin{bmatrix} 7 & 5 \\ 11 & 2 \end{bmatrix}$					
c) $\begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix}$	d) $\begin{bmatrix} 7 & -5 \\ 11 & 2 \end{bmatrix}$					
16.	The points on the curve $\frac{x^2}{9} + \frac{y^2}{16} = 1$ at which the tangents are parallel to y-axis are:	1				
<table><tr><td>a) $(0, \pm 4)$</td><td>b) $(\pm 4, 0)$</td></tr><tr><td>c) $(\pm 3, 0)$</td><td>d) $(0, \pm 3)$</td></tr></table>			a) $(0, \pm 4)$	b) $(\pm 4, 0)$	c) $(\pm 3, 0)$	d) $(0, \pm 3)$
a) $(0, \pm 4)$	b) $(\pm 4, 0)$					
c) $(\pm 3, 0)$	d) $(0, \pm 3)$					
17.	Given that $A = [a_{ij}]$ is a square matrix of order 3×3 and $ A = -7$, then the value of $\sum_{i=1}^3 a_{i2} A_{i2}$, where A_{ij} denotes the cofactor of element a_{ij} is:	1				
<table><tr><td>a) 7</td><td>b) -7</td></tr><tr><td>c) 0</td><td>d) 49</td></tr></table>			a) 7	b) -7	c) 0	d) 49
a) 7	b) -7					
c) 0	d) 49					
18.	If $y = \log(\cos e^x)$, then $\frac{dy}{dx}$ is:	1				
<table><tr><td>a) $\cos e^{x-1}$</td><td>b) $e^{-x} \cos e^x$</td></tr><tr><td>c) $e^x \sin e^x$</td><td>d) $-e^x \tan e^x$</td></tr></table>			a) $\cos e^{x-1}$	b) $e^{-x} \cos e^x$	c) $e^x \sin e^x$	d) $-e^x \tan e^x$
a) $\cos e^{x-1}$	b) $e^{-x} \cos e^x$					
c) $e^x \sin e^x$	d) $-e^x \tan e^x$					
19.	Based on the given shaded region as the feasible region in the graph, at which point(s) is the objective function $Z = 3x + 9y$ maximum?	1				
						
<table><tr><td>a) Point B</td><td>b) Point C</td></tr><tr><td>c) Point D</td><td>d) every point on the line segment CD</td></tr></table>			a) Point B	b) Point C	c) Point D	d) every point on the line segment CD
a) Point B	b) Point C					
c) Point D	d) every point on the line segment CD					


20.	The least value of the function $f(x) = 2\cos x + x$ in the closed interval $[0, \frac{\pi}{2}]$ is:	1				
	<table><tr><td>a) 2</td><td>b) $\frac{\pi}{6} + \sqrt{3}$</td></tr><tr><td>c) $\frac{\pi}{2}$</td><td>d) The least value does not exist.</td></tr></table>	a) 2	b) $\frac{\pi}{6} + \sqrt{3}$	c) $\frac{\pi}{2}$	d) The least value does not exist.	
a) 2	b) $\frac{\pi}{6} + \sqrt{3}$					
c) $\frac{\pi}{2}$	d) The least value does not exist.					

SECTION – B

**In this section, attempt any 16 questions out of the Questions 21 - 40.
Each Question is of 1 mark weightage.**

21.	The function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined as $f(x) = x^3$ is:	1				
	<table><tr><td>a) One-on but not onto</td><td>b) Not one-one but onto</td></tr><tr><td>c) Neither one-one nor onto</td><td>d) One-one and onto</td></tr></table>	a) One-on but not onto	b) Not one-one but onto	c) Neither one-one nor onto	d) One-one and onto	
a) One-on but not onto	b) Not one-one but onto					
c) Neither one-one nor onto	d) One-one and onto					

22.	If $x = a \sec \theta$, $y = b \tan \theta$, then $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{6}$ is:	1				
	<table><tr><td>a) $\frac{-3\sqrt{3}b}{a^2}$</td><td>b) $\frac{-2\sqrt{3}b}{a}$</td></tr><tr><td>c) $\frac{-3\sqrt{3}b}{a}$</td><td>d) $\frac{-b}{3\sqrt{3}a^2}$</td></tr></table>	a) $\frac{-3\sqrt{3}b}{a^2}$	b) $\frac{-2\sqrt{3}b}{a}$	c) $\frac{-3\sqrt{3}b}{a}$	d) $\frac{-b}{3\sqrt{3}a^2}$	
a) $\frac{-3\sqrt{3}b}{a^2}$	b) $\frac{-2\sqrt{3}b}{a}$					
c) $\frac{-3\sqrt{3}b}{a}$	d) $\frac{-b}{3\sqrt{3}a^2}$					

23.		In the given graph, the feasible region for a LPP is shaded. The objective function $Z = 2x - 3y$, will be minimum at:	1				
	<table><tr><td>a) (4, 10)</td><td>b) (6, 8)</td></tr><tr><td>c) (0, 8)</td><td>d) (6, 5)</td></tr></table>	a) (4, 10)	b) (6, 8)	c) (0, 8)	d) (6, 5)		
a) (4, 10)	b) (6, 8)						
c) (0, 8)	d) (6, 5)						

24.	The derivative of $\sin^{-1}(2x\sqrt{1-x^2})$ w.r.t $\sin^{-1}x$, $-\frac{1}{\sqrt{2}} < x < \frac{1}{\sqrt{2}}$, is:	1				
	<table><tr><td>a) 2</td><td>b) $\frac{\pi}{2} - 2$</td></tr><tr><td>c) $\frac{\pi}{2}$</td><td>d) -2</td></tr></table>	a) 2	b) $\frac{\pi}{2} - 2$	c) $\frac{\pi}{2}$	d) -2	
a) 2	b) $\frac{\pi}{2} - 2$					
c) $\frac{\pi}{2}$	d) -2					

25.	If $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -4 \\ 2 & -1 & 5 \end{bmatrix}$, then:	1				
	<table><tr><td>a) $A^{-1} = B$</td><td>b) $A^{-1} = 6B$</td></tr><tr><td>c) $B^{-1} = B$</td><td>d) $B^{-1} = \frac{1}{6}A$</td></tr></table>	a) $A^{-1} = B$	b) $A^{-1} = 6B$	c) $B^{-1} = B$	d) $B^{-1} = \frac{1}{6}A$	
a) $A^{-1} = B$	b) $A^{-1} = 6B$					
c) $B^{-1} = B$	d) $B^{-1} = \frac{1}{6}A$					

26.	The real function $f(x) = 2x^3 - 3x^2 - 36x + 7$ is:	1								
<table><tr><td colspan="2">a) Strictly increasing in $(-\infty, -2)$ and strictly decreasing in $(-2, \infty)$</td></tr><tr><td colspan="2">b) Strictly decreasing in $(-2, 3)$</td></tr><tr><td colspan="2">c) Strictly decreasing in $(-\infty, 3)$ and strictly increasing in $(3, \infty)$</td></tr><tr><td colspan="2">d) Strictly decreasing in $(-\infty, -2) \cup (3, \infty)$</td></tr></table>			a) Strictly increasing in $(-\infty, -2)$ and strictly decreasing in $(-2, \infty)$		b) Strictly decreasing in $(-2, 3)$		c) Strictly decreasing in $(-\infty, 3)$ and strictly increasing in $(3, \infty)$		d) Strictly decreasing in $(-\infty, -2) \cup (3, \infty)$	
a) Strictly increasing in $(-\infty, -2)$ and strictly decreasing in $(-2, \infty)$										
b) Strictly decreasing in $(-2, 3)$										
c) Strictly decreasing in $(-\infty, 3)$ and strictly increasing in $(3, \infty)$										
d) Strictly decreasing in $(-\infty, -2) \cup (3, \infty)$										
27.	Simplest form of $\tan^{-1} \left(\frac{\sqrt{1+\cos x} + \sqrt{1-\cos x}}{\sqrt{1+\cos x} - \sqrt{1-\cos x}} \right), \pi < x < \frac{3\pi}{2}$ is:	1								
<table><tr><td>a) $\frac{\pi}{4} - \frac{x}{2}$</td><td>b) $\frac{3\pi}{2} - \frac{x}{2}$</td></tr><tr><td>c) $-\frac{x}{2}$</td><td>d) $\pi - \frac{x}{2}$</td></tr></table>			a) $\frac{\pi}{4} - \frac{x}{2}$	b) $\frac{3\pi}{2} - \frac{x}{2}$	c) $-\frac{x}{2}$	d) $\pi - \frac{x}{2}$				
a) $\frac{\pi}{4} - \frac{x}{2}$	b) $\frac{3\pi}{2} - \frac{x}{2}$									
c) $-\frac{x}{2}$	d) $\pi - \frac{x}{2}$									
28.	Given that A is a non-singular matrix of order 3 such that $A^2 = 2A$, then value of $ 2A $ is:	1								
<table><tr><td>a) 4</td><td>b) 8</td></tr><tr><td>c) 64</td><td>d) 16</td></tr></table>			a) 4	b) 8	c) 64	d) 16				
a) 4	b) 8									
c) 64	d) 16									
29.	The value of b for which the function $f(x) = x + \cos x + b$ is strictly decreasing over \mathbf{R} is:	1								
<table><tr><td>a) $b < 1$</td><td>b) No value of b exists</td></tr><tr><td>c) $b \leq 1$</td><td>d) $b \geq 1$</td></tr></table>			a) $b < 1$	b) No value of b exists	c) $b \leq 1$	d) $b \geq 1$				
a) $b < 1$	b) No value of b exists									
c) $b \leq 1$	d) $b \geq 1$									
30.	Let R be the relation in the set N given by $R = \{(a, b) : a = b - 2, b > 6\}$, then:	1								
<table><tr><td>a) $(2, 4) \in R$</td><td>b) $(3, 8) \in R$</td></tr><tr><td>c) $(6, 8) \in R$</td><td>d) $(8, 7) \in R$</td></tr></table>			a) $(2, 4) \in R$	b) $(3, 8) \in R$	c) $(6, 8) \in R$	d) $(8, 7) \in R$				
a) $(2, 4) \in R$	b) $(3, 8) \in R$									
c) $(6, 8) \in R$	d) $(8, 7) \in R$									
31.	The point(s), at which the function f given by $f(x) = \begin{cases} \frac{x}{ x }, & x < 0 \\ -1, & x \geq 0 \end{cases}$ is continuous, is/are:	1								
<table><tr><td>a) $x \in \mathbf{R}$</td><td>b) $x = 0$</td></tr><tr><td>c) $x \in \mathbf{R} - \{0\}$</td><td>d) $x = -1$ and 1</td></tr></table>			a) $x \in \mathbf{R}$	b) $x = 0$	c) $x \in \mathbf{R} - \{0\}$	d) $x = -1$ and 1				
a) $x \in \mathbf{R}$	b) $x = 0$									
c) $x \in \mathbf{R} - \{0\}$	d) $x = -1$ and 1									
32.	If $A = \begin{bmatrix} 0 & 2 \\ 3 & -4 \end{bmatrix}$ and $kA = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$, then the values of k, a and b respectively are:	1								

	<table><tr><td>a) $-6, -12, -18$</td><td>b) $-6, -4, -9$</td></tr><tr><td>c) $-6, 4, 9$</td><td>d) $-6, 12, 18$</td></tr></table>	a) $-6, -12, -18$	b) $-6, -4, -9$	c) $-6, 4, 9$	d) $-6, 12, 18$	
a) $-6, -12, -18$	b) $-6, -4, -9$					
c) $-6, 4, 9$	d) $-6, 12, 18$					
33.	<p>A linear programming problem is as follows: <i>Minimize</i> $Z = 30x + 50y$ subject to the constraints, $3x + 5y \geq 15$ $2x + 3y \leq 18$ $x \geq 0, y \geq 0$ In the feasible region, the minimum value of Z occurs at</p> <table><tr><td>a) a unique point</td><td>b) no point</td></tr><tr><td>c) infinitely many points</td><td>d) two points only</td></tr></table>	a) a unique point	b) no point	c) infinitely many points	d) two points only	1
a) a unique point	b) no point					
c) infinitely many points	d) two points only					
34.	<p>The area of a trapezium is defined by function f and given by $f(x) = (10 + x)\sqrt{100 - x^2}$, then the area when it is maximised is:</p> <table><tr><td>a) 75cm^2</td><td>b) $7\sqrt{3}\text{cm}^2$</td></tr><tr><td>c) $75\sqrt{3}\text{cm}^2$</td><td>d) 5cm^2</td></tr></table>	a) 75cm^2	b) $7\sqrt{3}\text{cm}^2$	c) $75\sqrt{3}\text{cm}^2$	d) 5cm^2	1
a) 75cm^2	b) $7\sqrt{3}\text{cm}^2$					
c) $75\sqrt{3}\text{cm}^2$	d) 5cm^2					
35.	<p>If A is square matrix such that $A^2 = A$, then $(I + A)^3 - 7A$ is equal to:</p> <table><tr><td>a) A</td><td>b) $I + A$</td></tr><tr><td>c) $I - A$</td><td>d) I</td></tr></table>	a) A	b) $I + A$	c) $I - A$	d) I	1
a) A	b) $I + A$					
c) $I - A$	d) I					
36.	<p>If $\tan^{-1} x = y$, then:</p> <table><tr><td>a) $-1 < y < 1$</td><td>b) $\frac{-\pi}{2} \leq y \leq \frac{\pi}{2}$</td></tr><tr><td>c) $\frac{-\pi}{2} < y < \frac{\pi}{2}$</td><td>d) $y \in \{\frac{-\pi}{2}, \frac{\pi}{2}\}$</td></tr></table>	a) $-1 < y < 1$	b) $\frac{-\pi}{2} \leq y \leq \frac{\pi}{2}$	c) $\frac{-\pi}{2} < y < \frac{\pi}{2}$	d) $y \in \{\frac{-\pi}{2}, \frac{\pi}{2}\}$	1
a) $-1 < y < 1$	b) $\frac{-\pi}{2} \leq y \leq \frac{\pi}{2}$					
c) $\frac{-\pi}{2} < y < \frac{\pi}{2}$	d) $y \in \{\frac{-\pi}{2}, \frac{\pi}{2}\}$					
37.	<p>Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and let $f = \{(1, 4), (2, 5), (3, 6)\}$ be a function from A to B. Based on the given information, f is best defined as:</p> <table><tr><td>a) Surjective function</td><td>b) Injective function</td></tr><tr><td>c) Bijective function</td><td>d) function</td></tr></table>	a) Surjective function	b) Injective function	c) Bijective function	d) function	1
a) Surjective function	b) Injective function					
c) Bijective function	d) function					
38.	<p>For $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, then $14A^{-1}$ is given by:</p> <table><tr><td>a) $14 \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$</td><td>b) $\begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$</td></tr><tr><td>c) $2 \begin{bmatrix} 2 & -1 \\ 1 & -3 \end{bmatrix}$</td><td>d) $2 \begin{bmatrix} -3 & -1 \\ 1 & -2 \end{bmatrix}$</td></tr></table>	a) $14 \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$	b) $\begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$	c) $2 \begin{bmatrix} 2 & -1 \\ 1 & -3 \end{bmatrix}$	d) $2 \begin{bmatrix} -3 & -1 \\ 1 & -2 \end{bmatrix}$	1
a) $14 \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$	b) $\begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$					
c) $2 \begin{bmatrix} 2 & -1 \\ 1 & -3 \end{bmatrix}$	d) $2 \begin{bmatrix} -3 & -1 \\ 1 & -2 \end{bmatrix}$					
39.	<p>The point(s) on the curve $y = x^3 - 11x + 5$ at which the tangent is $y = x - 11$ is/are:</p> <table><tr><td>a) $(-2, 19)$</td><td>b) $(2, -9)$</td></tr><tr><td>c) $(\pm 2, 19)$</td><td>d) $(-2, 19)$ and $(2, -9)$</td></tr></table>	a) $(-2, 19)$	b) $(2, -9)$	c) $(\pm 2, 19)$	d) $(-2, 19)$ and $(2, -9)$	1
a) $(-2, 19)$	b) $(2, -9)$					
c) $(\pm 2, 19)$	d) $(-2, 19)$ and $(2, -9)$					
40.	<p>Given that $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ and $A^2 = 3I$, then:</p>	1				

a) $1 + \alpha^2 + \beta\gamma = 0$	b) $1 - \alpha^2 - \beta\gamma = 0$
c) $3 - \alpha^2 - \beta\gamma = 0$	d) $3 + \alpha^2 + \beta\gamma = 0$

SECTION – C

In this section, attempt any 8 questions.

Each question is of 1-mark weightage.

Questions 46-50 are based on a Case-Study.

41.	For an objective function $Z = ax + by$, where $a, b > 0$; the corner points of the feasible region determined by a set of constraints (linear inequalities) are $(0, 20)$, $(10, 10)$, $(30, 30)$ and $(0, 40)$. The condition on a and b such that the maximum Z occurs at both the points $(30, 30)$ and $(0, 40)$ is:	1				
	<table><tr><td>a) $b - 3a = 0$</td><td>b) $a = 3b$</td></tr><tr><td>c) $a + 2b = 0$</td><td>d) $2a - b = 0$</td></tr></table>	a) $b - 3a = 0$	b) $a = 3b$	c) $a + 2b = 0$	d) $2a - b = 0$	
a) $b - 3a = 0$	b) $a = 3b$					
c) $a + 2b = 0$	d) $2a - b = 0$					
42.	For which value of m is the line $y = mx + 1$ a tangent to the curve $y^2 = 4x$?	1				
	<table><tr><td>a) $\frac{1}{2}$</td><td>b) 1</td></tr><tr><td>c) 2</td><td>d) 3</td></tr></table>	a) $\frac{1}{2}$	b) 1	c) 2	d) 3	
a) $\frac{1}{2}$	b) 1					
c) 2	d) 3					
43.	The maximum value of $[x(x - 1) + 1]^{\frac{1}{3}}$, $0 \leq x \leq 1$ is:	1				
	<table><tr><td>a) 0</td><td>b) $\frac{1}{2}$</td></tr><tr><td>c) 1</td><td>d) $\sqrt[3]{\frac{1}{3}}$</td></tr></table>	a) 0	b) $\frac{1}{2}$	c) 1	d) $\sqrt[3]{\frac{1}{3}}$	
a) 0	b) $\frac{1}{2}$					
c) 1	d) $\sqrt[3]{\frac{1}{3}}$					
44.	In a linear programming problem, the constraints on the decision variables x and y are $x - 3y \geq 0, y \geq 0, 0 \leq x \leq 3$. The feasible region	1				
	<table><tr><td>a) is not in the first quadrant</td><td>b) is bounded in the first quadrant</td></tr><tr><td>c) is unbounded in the first quadrant</td><td>d) does not exist</td></tr></table>	a) is not in the first quadrant	b) is bounded in the first quadrant	c) is unbounded in the first quadrant	d) does not exist	
a) is not in the first quadrant	b) is bounded in the first quadrant					
c) is unbounded in the first quadrant	d) does not exist					
45.	Let $A = \begin{bmatrix} 1 & \sin\alpha & 1 \\ -\sin\alpha & 1 & \sin\alpha \\ -1 & -\sin\alpha & 1 \end{bmatrix}$, where $0 \leq \alpha \leq 2\pi$, then:	1				
	<table><tr><td>a) $A =0$</td><td>b) $A \in (2, \infty)$</td></tr><tr><td>c) $A \in (2, 4)$</td><td>d) $A \in [2, 4]$</td></tr></table>	a) $ A =0$	b) $ A \in (2, \infty)$	c) $ A \in (2, 4)$	d) $ A \in [2, 4]$	
a) $ A =0$	b) $ A \in (2, \infty)$					
c) $ A \in (2, 4)$	d) $ A \in [2, 4]$					

CASE STUDY



The fuel cost per hour for running a train is proportional to the square of the speed it generates in km per hour. If the fuel costs ₹ 48 per hour at speed 16 km per hour and the fixed charges to run the train amount to ₹ 1200 per hour.

Assume the speed of the train as v km/h.

	Based on the given information, answer the following questions.					
46.	Given that the fuel cost per hour is k times the square of the speed the train generates in km/h, the value of k is:	1				
	<table><tr><td>a) $\frac{16}{3}$</td><td>b) $\frac{1}{3}$</td></tr><tr><td>c) 3</td><td>d) $\frac{3}{16}$</td></tr></table>	a) $\frac{16}{3}$	b) $\frac{1}{3}$	c) 3	d) $\frac{3}{16}$	
a) $\frac{16}{3}$	b) $\frac{1}{3}$					
c) 3	d) $\frac{3}{16}$					
47.	If the train has travelled a distance of 500km, then the total cost of running the train is given by function:	1				
	<table><tr><td>a) $\frac{15}{16}v + \frac{600000}{v}$</td><td>b) $\frac{375}{4}v + \frac{600000}{v}$</td></tr><tr><td>c) $\frac{5}{16}v^2 + \frac{150000}{v}$</td><td>d) $\frac{3}{16}v + \frac{6000}{v}$</td></tr></table>	a) $\frac{15}{16}v + \frac{600000}{v}$	b) $\frac{375}{4}v + \frac{600000}{v}$	c) $\frac{5}{16}v^2 + \frac{150000}{v}$	d) $\frac{3}{16}v + \frac{6000}{v}$	
a) $\frac{15}{16}v + \frac{600000}{v}$	b) $\frac{375}{4}v + \frac{600000}{v}$					
c) $\frac{5}{16}v^2 + \frac{150000}{v}$	d) $\frac{3}{16}v + \frac{6000}{v}$					
48.	The most economical speed to run the train is:	1				
	<table><tr><td>a) 18km/h</td><td>b) 5km/h</td></tr><tr><td>c) 80km/h</td><td>d) 40km/h</td></tr></table>	a) 18km/h	b) 5km/h	c) 80km/h	d) 40km/h	
a) 18km/h	b) 5km/h					
c) 80km/h	d) 40km/h					
49.	The fuel cost for the train to travel 500km at the most economical speed is:	1				
	<table><tr><td>a) ₹ 3750</td><td>b) ₹ 750</td></tr><tr><td>c) ₹ 7500</td><td>d) ₹ 75000</td></tr></table>	a) ₹ 3750	b) ₹ 750	c) ₹ 7500	d) ₹ 75000	
a) ₹ 3750	b) ₹ 750					
c) ₹ 7500	d) ₹ 75000					
50.	The total cost of the train to travel 500km at the most economical speed is:	1				
	<table><tr><td>a) ₹ 3750</td><td>b) ₹ 75000</td></tr><tr><td>c) ₹ 7500</td><td>d) ₹ 15000</td></tr></table>	a) ₹ 3750	b) ₹ 75000	c) ₹ 7500	d) ₹ 15000	
a) ₹ 3750	b) ₹ 75000					
c) ₹ 7500	d) ₹ 15000					
