

Answer to Some Selected Problems

UNIT 7

7.25 15 g

UNIT 8

8.32 Mass of carbon dioxide formed = 0.505 g

Mass of water formed = 0.0864 g

8.33 % of nitrogen = 56

8.34 % of chlorine = 37.57

8.35 % of sulphur = 19.66

UNIT 9

9.1 Due to the side reaction in termination step by the combination of two $\dot{\text{C}}\text{H}_3$ free radicals.

- | | | |
|-----|--------------------------------|-------------------------------|
| 9.2 | (a) 2-Methyl-but-2-ene | (b) Pent-1-ene-3-yne |
| | (c) Buta-1, 3-diene | (d) 4-Phenylbut-1-ene |
| | (e) 2-Methylphenol | (f) 5-(2-Methylpropyl)-decane |
| | (g) 4-Ethyldeca -1,5,8- triene | |

- | | | |
|-----|--|-----------------|
| 9.3 | (a) (i) $\text{CH}_2 = \text{CH} - \text{CH}_2 - \text{CH}_3$ | But-1-ene |
| | (ii) $\text{CH}_3 - \text{CH}_2 = \text{CH} - \text{CH}_3$ | But-2-ene |
| | (iii) $\text{CH}_2 = \underset{\text{CH}_3}{\text{C}} - \text{CH}_3$ | 2-Methylpropene |


- | | | |
|---------|---|-------------------|
| (b) (i) | $\text{HC} \equiv \text{C} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$ | Pent-1-yne |
| (ii) | $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_2 - \text{CH}_3$ | Pent-2-yne |
| (iii) | $\text{CH}_3 - \text{CH} - \underset{\text{CH}_3}{\text{C}} \equiv \text{CH}$ | 3-Methylbut-1-yne |

- | | | |
|-----|---------------------------------|-----------------------------------|
| 9.4 | (i) Ethanal and propanal | (ii) Butan-2-one and pentan-2-one |
| | (iii) Methanal and pentan-3-one | (iv) Propanal and benzaldehyde |


9.5 3-Ethylpent-2-ene

9.6 But-2-ene

9.7 4-Ethylhex-3-ene





- 9.8 (a) $\text{C}_4\text{H}_{10}(\text{g}) + 13/20_2(\text{g}) \xrightarrow{\Delta} 4\text{CO}_2(\text{g}) + 5\text{H}_2\text{O}(\text{g})$
 (b) $\text{C}_5\text{H}_{10}(\text{g}) + 15/20_2(\text{g}) \xrightarrow{\Delta} 5\text{CO}_2(\text{g}) + 5\text{H}_2\text{O}(\text{g})$
 (c) $\text{C}_5\text{H}_{10}(\text{g}) + 17/2 \text{O}_2(\text{g}) \xrightarrow{\Delta} 6\text{CO}_2(\text{g}) + 5\text{H}_2\text{O}(\text{g})$
 (d) $\text{C}_7\text{H}_8(\text{g}) + 9\text{O}_2(\text{g}) \xrightarrow{\Delta} 7\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\text{g})$





The *cis* form will have higher boiling point due to more polar nature leading to stronger intermolecular dipole-dipole interaction, thus requiring more heat energy to separate them.


- 9.10 Due to resonance
 9.11 Planar, conjugated ring system with delocalisation of $(4n+2)$ electrons, where, n is an integer
 9.12 Lack of delocalisation of $(4n+2)$ π electrons in the cyclic system.
 9.13 (i)



(iii)



9.14




15 H attached to 1 carbons

4 H attached to 2 carbons

1 H attaches to 3 carbons

9.15 More the branching in alkane, lower will be the boiling point.

9.16 Refer to addition reaction of HBr to unsymmetrical alkenes in the text.



All the three products cannot be obtained by any one of the Kekulé's structures. This shows that benzene is a resonance hybrid of the two resonating structures.

9.18 $\text{H} - \text{C} \equiv \text{C} - \text{H} > \text{C}_6\text{H}_6 > \text{C}_6\text{H}_{14}$. Due to maximum s orbital character in ethyne (50 per cent) as compared to 33 per cent in benzene and 25 per cent in n-hexane.

9.19 Due to the presence of 6 π electrons, benzene behaves as a rich source of electrons thus being easily attacked by reagents deficient in electrons.

- 9.20 (i) $3 \text{ CH} \equiv \text{CH} \xrightarrow[873\text{K}]{\text{Red hot Iron tube}} \text{C}_6\text{H}_6$
- (ii) $\text{C}_2\text{H}_4 \xrightarrow{\text{Br}_2} \begin{array}{c} \text{CH}_2 - \text{CH}_2 \\ | \quad | \\ \text{Br} \quad \text{BR} \end{array} \xrightarrow{\text{alc. KOH}} \text{CH}_2 = \text{CHN}r \xrightarrow{\text{NaNH}_2} \text{HC} \equiv \text{HC} \xrightarrow[873\text{k}]{\text{Red hot Iron tube}} \text{C}_6\text{H}_6$
- (iii) $\text{C}_6\text{H}_{14} \xrightarrow[773 \text{ K, 10-20 atm}]{\text{Cr}_2\text{O}_3 / \text{V}_2\text{O}_5 / \text{Mo}_2\text{O}_3} \text{C}_6\text{H}_6$
- 9.21 $\text{CH}_2 = \text{C}(\text{CH}_3) - \text{CH}_2 - \text{CH}_3$ 2-Methylbut-1-ene
- $\text{CH}_3 - \text{C}(\text{CH}_3) = \text{CH} - \text{CH}_3$ 2-Methylbut-2-ene
- $\text{CH}_3 - \text{CH}(\text{CH}_3) - \text{CH} = \text{CH}_2$ 3-Methylbut-1-ene
- 9.22 (a) Chlorobenzene > *p*-nitrochlorobenzene > 2,4-dinitrochlorobenzene
 (b) Toluene > *p*-CH₃-C₆H₄-NO₂ > *p*-O₂N-C₆H₄-NO₂
- 9.23 Toluene undergoes nitration most easily due to electron releasing nature of the methyl group.
- 9.24 FeCl₃
- 9.25 Due to the formation of side products. For example, by starting with 1-bromopropane and 1-bromobutane, hexane and octane are the side products besides heptane.

NOTES

© NCERT
not to be republished