


Class 8 Maths Ganita Prakash Chapter 4 Quadrilaterals NCERT Solutions


Textbook Page 94

Figure it Out

1. Find all the other angles inside the following rectangles.




(i)



(ii)

Solution:

(i)



$\angle 1 + \angle 9 = 90^\circ$ (All corner angles of a rectangle are 90°)

$$\angle 1 + 30^\circ = 90^\circ$$

$$\angle 1 = 90^\circ - 30^\circ$$

$$\angle 1 = 60^\circ \quad \angle 1 = \angle 5 = 60^\circ \text{ (Alternate interior angles)}$$

$$\angle 9 = \angle 4 = 30^\circ \text{ (Alternate interior angles)}$$

In $\triangle AOB$, $OA = OB$, then the angles opposite them are equal

$$\therefore \angle 9 = \angle 7 = 30^\circ \quad \angle 7 = \angle 3 = 30^\circ \text{ (Alternate interior angles)}$$

In $\triangle AOD$, $OA = OD$, then the angles opposite them are equal

$$\therefore \angle 2 = \angle 1 = 60^\circ \quad \angle 2 = \angle 6 = 60^\circ \text{ (Alternate interior angles)}$$

In $\triangle AOB$,

$$\angle 9 + \angle 7 + \angle AOB = 180^\circ \text{ (Sum of angles of a triangle)}$$

$$30^\circ + 30^\circ + \angle AOB = 180^\circ$$

$$60^\circ + \angle AOB = 180^\circ$$

$$\angle AOB = 180^\circ - 60^\circ$$

$$\angle AOB = 120^\circ$$

$$\angle AOB = \angle COD = 120^\circ \text{ (Vertically opposite angles)}$$

$\angle AOB + \angle AOD = 180^\circ$ (Linear pair)
 $120^\circ + \angle AOD = 180^\circ$
 $\angle AOD = 180^\circ - 120^\circ$
 $\angle AOD = 60^\circ$


$\angle AOD = \angle BOC = 60^\circ$ (Vertically opposite angles)

Therefore, $\angle 1 = \angle 5 = \angle 2 = \angle 6 = \angle AOD = \angle BOC = 60^\circ$.

$\angle AOB = \angle COD = 120^\circ$.

$\angle 9 = \angle 4 = \angle 7 = \angle 3 = 30^\circ$.

(ii)



$\angle POS = \angle ROQ = 110^\circ$ (Vertically opposite angles)

$\angle POS + \angle POQ = 180^\circ$ (Linear Pair)

$110^\circ + \angle POQ = 180^\circ$

$\angle POQ = 180^\circ - 110^\circ$

$\angle POQ = 70^\circ$

$\angle POQ = \angle SOR = 70^\circ$ (Vertically opposite angles)

In $\triangle POS$, $OP = OS$, then the angles opposite them are equal.

$\therefore \angle 1 = \angle 2 = a$ In $\triangle POS$,

$\angle 1 + \angle 2 + \angle POS = 180^\circ$ (Sum of angles of a triangle)

$a + a + 110^\circ = 180^\circ$

$2a = 180^\circ - 110^\circ$

$2a = 70^\circ$

$a = 35^\circ$

$\therefore \angle 1 = \angle 2 = a = 35^\circ$

$$\angle 1 = \angle 5 = 35^\circ \dots\dots\dots (\text{Alternate interior angles})$$

$$\angle 2 = \angle 6 = 35^\circ \dots\dots\dots (\text{Alternate interior angles})$$

Since ABCD is a rectangle, $\angle P = 90^\circ$

$$\angle 9 = \angle 1 + \angle 8$$

$$90^\circ = 35^\circ + \angle 8$$

$$\angle 8 = 90^\circ - 35^\circ$$

$$\angle 8 = 55^\circ$$

$$\angle 8 = \angle 4 = 55^\circ \dots\dots\dots (\text{Alternate interior angles})$$

In $\triangle POQ$, $OP = OQ$, then the angles opposite them are equal

i.e. $\angle 7 = \angle 8 = 55^\circ$

$$\angle 7 = \angle 2 = 55^\circ \dots\dots\dots (\text{Alternate interior angles})$$

Therefore, $\angle POS = \angle ROQ = 110^\circ$.

$$\angle POQ = \angle SOR = 70^\circ$$

$$\angle 1 = \angle 2 = \angle 5 = \angle 6 = 35^\circ$$


$$\angle 8 = \angle 4 = \angle 7 = \angle 2 = 55^\circ$$

2. Draw a quadrilateral whose diagonals have equal lengths of 8 cm that bisect each other, and intersect at an angle of


- (i) 30° (ii) 40° (iii) 90° (iv) 140°**

Solution:


- (i) 30°**




(ii) 40°



(iii) 90°




(iv) 140°



3. Consider a circle with centre O. Line segments PL and AM are two perpendicular diameters of the circle. What is the figure APML? Reason and/or experiment to figure this out.

Solution:



Let PL and AM be two perpendicular diameters of a circle with centre O and radius r .

We have,

$$PO = OL = AO = OM = r$$

$$\therefore PL = PO + OL = r + r = 2r$$

$$\text{and } AM = AO + OM = r + r = 2r$$

Hence, $PL = AM$.

In quadrilateral APML, the diagonals PL and AM are equal in length and are perpendicular bisectors of each other.

\therefore APML is a square.

4. We have seen how to get 90° using paper folding. Now, suppose we do not have any paper but two sticks of equal length and a thread. How do we make an exact 90° using these?

Solution:

Arrange the two sticks to cross each other and tie their ends with a thread to form a rhombus. Adjust the thread so all sides are equal. The diagonals of a rhombus intersect at 90° , so the sticks will form an exact right angle.





5. We saw that one of the properties of a rectangle is that its opposite sides are parallel. Can this be chosen as a definition of a rectangle? In other words, is every quadrilateral that has opposite sides parallel and equal a rectangle?

Solution:

No, this can't be the definition of a rectangle. A quadrilateral with opposite sides parallel and equal is a parallelogram, but not all parallelograms are rectangles. A rectangle needs all angles to be right angles.

Textbook Page 102

1. Find the remaining angles in the following quadrilaterals.



Solution:

(i) Here $PR \parallel EA$, and $PE \parallel RA$

Therefore, PEAR is a parallelogram.

$\angle P = \angle A = 40^\circ$ (Opposite angles of a parallelogram are equal)

$\angle P + \angle R = 180^\circ$ (The sum of the adjacent angles of a parallelogram is 180°)

$40^\circ + \angle R = 180^\circ$

$\angle R = 180^\circ - 40^\circ$

$\angle R = 140^\circ$.

$\angle R = \angle E = 140^\circ$ (Opposite angles of a parallelogram are equal)

(ii) Here $PQ \parallel SR$, and $PS \parallel QR$

\therefore PQRS is a parallelogram.

$\angle P = \angle R = 110^\circ$ (Opposite angles of a parallelogram are equal)

$\angle P + \angle S = 180^\circ$ (The sum of the adjacent angles of a parallelogram is 180°)
 $110^\circ + \angle S = 180^\circ$
 $\angle S = 180^\circ - 110^\circ$
 $\angle S = 70^\circ$.
 $\angle S = \angle Q = 70^\circ$ (Opposite angles of a parallelogram are equal)

(iii) Here, XWUV is a rhombus (all sides equal).

In $\triangle VUX$, $UV = UX$, then the angles opposite them are equal.

$\therefore \angle UXV = \angle UVX = 30^\circ$

$\angle UXV = \angle WXV = 30^\circ$ (The diagonals of a rhombus bisect its angles)

Also, $\angle UVX = \angle WVX = 30^\circ$ (The diagonals of a rhombus bisect its angles) $\angle E = 2 \times$

$\angle UVX = 2 \times 30^\circ = 60^\circ$

$\angle V = \angle X = 60^\circ$ (Opposite angles of a rhombus are equal)

$\angle V + \angle U = 180^\circ$ (The sum of adjacent angles of a rhombus is 180°)

$60^\circ + \angle U = 180^\circ$

$\angle U = 180^\circ - 60^\circ$

$\angle U = 120^\circ$

$\angle U = \angle W = 120^\circ$ (Opposite angles of a rhombus are equal)

(iv) Here, AEIO is a rhombus (all sides equal).

In $\triangle EAO$, $AE = AO$, then the angles opposite them are equal.

$\therefore \angle AOE = \angle AEO = 20^\circ$

$\angle AEO = \angle IEO = 20^\circ$ (The diagonals of a rhombus bisect its angles)

Also, $\angle AOE = \angle IOE = 20^\circ$ (The diagonals of a rhombus bisect its angles) $\angle E = 2 \times$

$\angle AEO = 2 \times 20^\circ = 40^\circ$

$\angle E = \angle O = 40^\circ$ (Opposite angles of a rhombus are equal)

$\angle E + \angle A = 180^\circ$ (The sum of adjacent angles of a rhombus is 180°)

$40^\circ + \angle A = 180^\circ$


$\angle A = 180^\circ - 40^\circ$

$\angle A = 140^\circ$

$\angle A = \angle I = 140^\circ$ (Opposite angles of a rhombus are equal)

2. Using the diagonal properties, construct a parallelogram whose diagonals are of lengths 7 cm and 5 cm, and intersect at an angle of 140° .

Solution:




Steps of construction:

- (i) Draw a line segment AC of length 7 cm and mark its midpoint as O.
- (ii) At point O, draw an angle of 140° with respect to diagonal AC.
- (iii) From O, along the 140° line in both directions, mark $OD = 2.5$ cm and $OB = 2.5$ cm using a compass.
- (iv) Join D to A and C.
Join B to A and C.
ABCD is the required parallelogram.

3. Using the diagonal properties, construct a rhombus whose diagonals are of lengths 4 cm and 5 cm.

Solution:




Steps of construction:

- (i) Draw a line segment AC of length 5 cm.
- (ii) Draw the perpendicular bisector of AC, intersecting it at O.
- (iii) With O as centre and radius 2 cm, mark points B (below) and D (above) on the perpendicular bisector.
- (iv) Join A–D, D–C, C–B, and B–A.
ABCD is the required rhombus.

Textbook Page 107

1. Find all the sides and the angles of the quadrilateral obtained by joining two equilateral triangles with sides 4 cm.

Solution:



Since all sides of an equilateral triangle are equal.

Thus, the lengths of all sides of the given quadrilateral are equal.

$\therefore PQ = QR = RS = SP = 4 \text{ cm}$.

Also, the measure of all angles of an equilateral triangle is 60° .


$\angle P = \angle R = 60^\circ$

$\angle S = \angle PSQ + \angle RSQ = 60^\circ + 60^\circ = 120^\circ$.

$\angle Q = \angle PQR + \angle RQS = 60^\circ + 60^\circ = 120^\circ$.


2. Construct a kite whose diagonals are of lengths 6 cm and 8 cm.

Solution:




- (i) Draw a line segment $AC = 6$ cm.
 - (ii) Construct the perpendicular bisector of AC ; let it meet AC at O (so O is the midpoint).
 - (iii) With centre O and radius 3 cm draw an arc to cut the bisector above AC ; label that point D . With centre O and radius 5 cm draw an arc to cut the bisector below AC ; label that point B .
 - (iv) Join $A - B$, $B - C$, $C - D$, $D - A$.
- $ABCD$ is the required kite.

3. Find the remaining angles in the following trapeziums —



Solution:



Since $AB \parallel DC$, and AD is a transversal, then

$\angle A + \angle D = 180^\circ$ (Sum of angles on the same side of the transversal)

$$135^\circ + \angle D = 180^\circ$$

$$\angle D = 180^\circ - 135^\circ$$

$$\angle D = 45^\circ$$


Also, since $AB \parallel DC$, and BC is a transversal, then

$\angle B + \angle C = 180^\circ$ (Sum of angles on the same side of the transversal)

$$105^\circ + \angle C = 180^\circ$$

$$\angle C = 180^\circ - 105^\circ$$

$$\angle C = 75^\circ$$



Since $PQ \parallel SR$, and PS is a transversal, then

$\angle P + \angle S = 180^\circ$ (Sum of angles on the same side of the transversal)

$$\angle P + 100^\circ = 180^\circ$$

$$\angle P = 180^\circ - 100^\circ = 80^\circ.$$

$\angle S = \angle R = 100^\circ$ (Angles opposite to equal sides are equal)

Also, since $PQ \parallel SR$, and QR is a transversal, then

$\angle Q + \angle R = 180^\circ$ (Sum of angles on the same side of the transversal)

$$\angle Q + 100^\circ = 180^\circ$$

$$\angle Q = 180^\circ - 100^\circ = 80^\circ.$$

4. Draw a Venn diagram showing the set of parallelograms, kites, rhombuses, rectangles, and squares. Then, answer the following questions —

(i) What is the quadrilateral that is both a kite and a parallelogram?

(ii) Can there be a quadrilateral that is both a kite and a rectangle?

(iii) Is every kite a rhombus? If not, what is the correct relationship between these two types of quadrilaterals?

Solution:


(i) A rhombus is a quadrilateral that is both a kite and a parallelogram.

(ii) A square is a quadrilateral that is both a kite and a rectangle.

(iii) No, every kite is not a rhombus.

Correct relationship: Every rhombus is a kite, but not every kite is a rhombus.

5. If PAIR and RODS are two rectangles, find $\angle IOD$.



Solution:

Since PAIR and RODS are two rectangles.

$\angle RIO = 90^\circ$ (Corner angle of a rectangle)

In $\triangle RIO$,

$\angle IRO + \angle IOR + \angle RIO = 180^\circ$ (Sum of angles of a triangle)

$$30^\circ + \angle IOR + 90^\circ = 180^\circ$$


$$120^\circ + \angle IOR = 180^\circ$$

$$\angle IOR = 180^\circ - 120^\circ = 60^\circ.$$

$$\therefore \angle IOD = 90^\circ - \angle IOR = 90^\circ - 60^\circ = 30^\circ.$$

6. Construct a square with a diagonal 6 cm without using a protractor.

Solution:



Steps of construction:


- (i) Draw a line segment $AC = 6$ cm and mark its midpoint as O.
- (ii) With O as centre and radius greater than half of AC, draw arcs above and below AC from points A and C.
- (iii) Join the arc intersections to get a line perpendicular to AC and passing through O.

(iv) Again, with O as centre and radius equal to 3 cm, mark points B and D on the perpendicular line.


(v) Connect A–B–C–D–A.

Hence, ABCD is the required square with a diagonal of 6 cm.

7. CASE is a square. The points U, V, W and X are the midpoints of the sides of the square. What type of quadrilateral is UVWX? Find this by using geometric reasoning, as well as by construction and measurement. Find other ways of constructing a square within a square such that the vertices of the inner square lie on the sides of the outer square, as shown in Figure (b).



(a)



(b)

Solution:

8. If a quadrilateral has four equal sides and one angle of 90° , will it be a square? Find the answer using geometric reasoning as well as by construction and measurement.

Solution:

Reasoning:

A rhombus is a quadrilateral with four equal sides.

If a rhombus has one angle of 90° , then:


Its opposite angle is also 90° (opposite angles of a rhombus are equal).

Each adjacent angle must also be 90° (sum of adjacent angles in a parallelogram/rhombus is 180°).

Thus, all four angles are 90° .

Since the quadrilateral has all sides equal and all angles right angles, it is a square.

Construction and measurement:



Steps of construction:

- (i) Draw a line segment PQ of length 5 cm.
- (ii) At point P, construct a perpendicular line to PQ.
- (iii) On this perpendicular, mark point S such that PS = 5 cm.
- (iv) With S as centre and radius 5 cm, draw an arc to the right of PS.
- (v) With Q as centre and radius 5 cm, draw an arc above PQ to intersect the arc from step (4) at point R.

Join Q–R, R–S, and S–P to complete the square PQRS. Verification by measurement:

All sides: $PQ = QR = RS = SP = 5 \text{ cm}$

All angles: $\angle P = \angle Q = \angle R = \angle S = 90^\circ$.

Conclusion: The figure constructed is a square.


9. What type of quadrilateral is one in which the opposite sides are equal? Justify your answer.

Hint: Draw a diagonal and check for congruent triangles.

Solution:

If a quadrilateral has opposite sides equal, then it is a parallelogram.

Geometric reasoning using a diagonal:



Given: Quadrilateral ABCD with $AB = CD$ and $BC = DA$.

Draw diagonal AC.

In $\triangle ABC$ and $\triangle CDA$,

$AB = CD$ (given)

$BC = DA$ (given)

$AC = AC$ (common side)

By SSS congruence, $\triangle ABC \cong \triangle CDA$.

From congruence, corresponding angles are equal:

$\angle BAC = \angle DCA$ and $\angle ACB = \angle CAD$.


But these are alternate interior angles.

$\therefore AB \parallel DC$ and $AD \parallel BC$.

Hence, ABCD is a parallelogram.

10. Will the sum of the angles in a quadrilateral such as the following one also be 360° ?


Find the answer using geometric reasoning as well as by constructing this figure and measuring.



Solution:

Yes, the sum of the angles in a quadrilateral will always be 360° .

Construction: Mark four non-collinear points as A, B, C, and D, and join them to form a quadrilateral ABCD.



Geometric reasoning:

In quad. ABCD, join BD to divide it into two triangles.

Now, In $\triangle BAD$,

$$\angle DBA + \angle BAD + \angle ADB = 180^\circ \dots\dots\dots(1)\dots\dots \text{(Sum of angles of a triangle)}$$

In $\triangle BCD$,

$$\angle BCD + \angle CDB + \angle DBC = 180^\circ \dots\dots\dots(2)\dots\dots \text{(Sum of angles of a triangle)}$$

Adding (1) and (2), we get

$$\angle DBA + \angle BAD + \angle ADB + \angle BCD + \angle CDB + \angle DBC = 180^\circ + 180^\circ$$

$$(\angle DBA + \angle DBC) + (\angle ADB + \angle CDB) + \angle BAD + \angle BCD = 360^\circ$$

$$\angle ABC + \angle ADC + \angle BAD + \angle BCD = 360^\circ$$

Thus, the sum of the angles of the given quadrilateral is 360° .

11. State whether the following statements are true or false. Justify your answers.

(i) A quadrilateral whose diagonals are equal and bisect each other must be a square.

Solution:

False.

A quadrilateral whose diagonals are equal and bisect each other is a rectangle. A square is a special case of a rectangle where all sides are also equal.

(ii) A quadrilateral having three right angles must be a rectangle.

Solution:

True.

Three right angles force the fourth to be right as well and a quadrilateral with four right angles is a rectangle.

(iii) A quadrilateral whose diagonals bisect each other must be a parallelogram.

Solution:

True.

If the diagonals bisect each other, then the two triangles formed by a diagonal are congruent, which gives pairs of opposite sides parallel. Hence the figure is a parallelogram.

(iv) A quadrilateral whose diagonals are perpendicular to each other must be a rhombus.

Solution:

False.

Squares, kites, and some other quadrilaterals also have perpendicular diagonals. Therefore, having perpendicular diagonals does not necessarily mean the quadrilateral is a rhombus.

(v) A quadrilateral in which the opposite angles are equal must be a parallelogram.

Solution:

True.

If both pairs of opposite angles are equal, then each pair of adjacent angles are supplementary, which implies opposite sides are parallel. Hence the quadrilateral is a parallelogram.

(vi) A quadrilateral in which all the angles are equal is a rectangle.

Solution:

True

If all four angles are equal, each angle must be $360^\circ/4 = 90^\circ$. A quadrilateral with four right angles is a rectangle.

(vii) Isosceles trapeziums are parallelograms.

Solution:

False.

An isosceles trapezium has exactly one pair of parallel sides and the non-parallel sides equal while a parallelogram must have two pairs of parallel sides. So an isosceles trapezium is not a parallelogram.